

平成29年度 事業年報

①人間ドック

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|-------|------|--------|---------------|--------|-----|-------|-------|--------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 0 | | 2 | 2.30% | 44 | 50.57% | 4 | 4.60% | 32 | 36.78% | 5 | 5.75% |
| | 女 | 95 | 0 | | 4 | 4.21% | 50 | 52.63% | 2 | 2.11% | 34 | 35.79% | 5 | 5.26% |
| | 計 | 182 | 0 | | 6 | 3.30% | 94 | 51.65% | 6 | 3.30% | 66 | 36.26% | 10 | 5.49% |
| 40 ～ 74 | 男 | 631 | 1 | 0.16% | 199 | 31.54% | 206 | 32.65% | 23 | 3.65% | 200 | 31.70% | 199 | 31.54% |
| | 女 | 635 | 0 | | 165 | 25.98% | 254 | 40.00% | 37 | 5.83% | 174 | 27.40% | 165 | 25.98% |
| | 計 | 1266 | 1 | 0.08% | 364 | 28.75% | 460 | 36.33% | 60 | 4.74% | 374 | 29.54% | 364 | 28.75% |
| 75 ～ | 男 | 22 | 0 | | 12 | 54.55% | 5 | 22.73% | 0 | | 5 | 22.73% | 12 | 54.55% |
| | 女 | 12 | 0 | | 9 | 75.00% | 1 | 8.33% | 1 | 8.33% | 1 | 8.33% | 9 | 75.00% |
| | 計 | 34 | 0 | | 21 | 61.76% | 6 | 17.65% | 1 | 2.94% | 6 | 17.65% | 21 | 61.76% |
| 合計 | | 1482 | 1 | 0.07% | 391 | 26.38% | 560 | 37.79% | 67 | 4.52% | 446 | 30.09% | 395 | 26.65% |

1. BMI

| 年代 | 性別 | 受診者数 | ～ 18.4 | | 18.5 ～ 24.9 | | 25.0 ～ | |
|---------|----|------|--------|--------|-------------|--------|--------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 5 | 5.75% | 50 | 57.47% | 32 | 36.78% |
| | 女 | 95 | 17 | 17.89% | 72 | 75.79% | 6 | 6.32% |
| | 計 | 182 | 22 | 12.09% | 122 | 67.03% | 38 | 20.88% |
| 40 ～ 74 | 男 | 631 | 13 | 2.06% | 424 | 67.19% | 194 | 30.74% |
| | 女 | 635 | 71 | 11.18% | 458 | 72.13% | 106 | 16.69% |
| | 計 | 1266 | 84 | 6.64% | 882 | 69.67% | 300 | 23.70% |
| 75 ～ | 男 | 22 | 2 | 9.09% | 16 | 72.73% | 4 | 18.18% |
| | 女 | 12 | 1 | 8.33% | 6 | 50.00% | 5 | 41.67% |
| | 計 | 34 | 3 | 8.82% | 22 | 64.71% | 9 | 26.47% |
| 合計 | | 1482 | 109 | 7.35% | 1026 | 69.23% | 347 | 23.41% |

2. 血圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|--------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 67 | 77.01% | 11 | 12.64% | 6 | 6.90% | 0 | | 2 | 2.30% | 1 | 1.15% |
| | 女 | 95 | 88 | 92.63% | 2 | 2.11% | 5 | 5.26% | 0 | | 0 | | 0 | |
| | 計 | 182 | 155 | 85.16% | 13 | 7.14% | 11 | 6.04% | 0 | | 2 | 1.10% | 1 | 0.55% |
| 40 ～ 74 | 男 | 631 | 285 | 45.17% | 93 | 14.74% | 101 | 16.01% | 0 | | 26 | 4.12% | 126 | 19.97% |
| | 女 | 635 | 386 | 60.79% | 79 | 12.44% | 72 | 11.34% | 0 | | 16 | 2.52% | 82 | 12.91% |
| | 計 | 1266 | 671 | 53.00% | 172 | 13.59% | 173 | 13.67% | 0 | | 42 | 3.32% | 208 | 16.43% |
| 75 ～ | 男 | 22 | 6 | 27.27% | 1 | 4.55% | 2 | 9.09% | 0 | | 4 | 18.18% | 9 | 40.91% |
| | 女 | 12 | 0 | | 1 | 8.33% | 3 | 25.00% | 0 | | 0 | | 8 | 66.67% |
| | 計 | 34 | 6 | 17.65% | 2 | 5.88% | 5 | 14.71% | 0 | | 4 | 11.76% | 17 | 50.00% |
| 合計 | | 1482 | 832 | 56.14% | 187 | 12.62% | 189 | 12.75% | 0 | | 48 | 3.24% | 226 | 15.25% |

3. 聴力検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 86 | 98.85% | 0 | 0 | 1 | 1.15% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 95 | 94 | 98.95% | 0 | 0 | 1 | 1.05% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 182 | 180 | 98.90% | 0 | 0 | 2 | 1.10% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 631 | 544 | 86.21% | 0 | 0 | 87 | 13.79% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 634 | 592 | 93.38% | 0 | 0 | 42 | 6.62% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 1265 | 1136 | 89.80% | 0 | 0 | 129 | 10.20% | 0 | 0 | 0 | 0 | 0 | 0 |
| 75 ～ | 男 | 22 | 13 | 59.09% | 0 | 0 | 9 | 40.91% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 12 | 6 | 50.00% | 0 | 0 | 6 | 50.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 34 | 19 | 55.88% | 0 | 0 | 15 | 44.12% | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 1481 | 1335 | 90.14% | 0 | 0 | 146 | 9.86% | 0 | 0 | 0 | 0 | 0 | 0 |

4. 肝機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 32 | 36.78% | 12 | 13.79% | 24 | 27.59% | 0 | 0 | 19 | 21.84% | 0 | 0 |
| | 女 | 95 | 75 | 78.95% | 3 | 3.16% | 12 | 12.63% | 0 | 0 | 5 | 5.26% | 0 | 0 |
| | 計 | 182 | 107 | 58.79% | 15 | 8.24% | 36 | 19.78% | 0 | 0 | 24 | 13.19% | 0 | 0 |
| 40 ～ 74 | 男 | 631 | 283 | 44.85% | 99 | 15.69% | 161 | 25.52% | 0 | 0 | 88 | 13.95% | 0 | 0 |
| | 女 | 635 | 460 | 72.44% | 50 | 7.87% | 87 | 13.70% | 0 | 0 | 36 | 5.67% | 2 | 0.31% |
| | 計 | 1266 | 743 | 58.69% | 149 | 11.77% | 248 | 19.59% | 0 | 0 | 124 | 9.79% | 2 | 0.16% |
| 75 ～ | 男 | 22 | 8 | 36.36% | 4 | 18.18% | 7 | 31.82% | 0 | 0 | 3 | 13.64% | 0 | 0 |
| | 女 | 12 | 7 | 58.33% | 3 | 25.00% | 1 | 8.33% | 0 | 0 | 1 | 8.33% | 0 | 0 |
| | 計 | 34 | 15 | 44.12% | 7 | 20.59% | 8 | 23.53% | 0 | 0 | 4 | 11.76% | 0 | 0 |
| 合計 | | 1482 | 865 | 58.37% | 171 | 11.54% | 292 | 19.70% | 0 | 0 | 152 | 10.26% | 2 | 0.13% |

5. 脂質検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 22 | 25.29% | 23 | 26.44% | 37 | 42.53% | 4 | 4.60% | 0 | 0 | 1 | 1.15% |
| | 女 | 95 | 57 | 60.00% | 10 | 10.53% | 21 | 22.11% | 2 | 2.11% | 5 | 5.26% | 0 | 0 |
| | 計 | 182 | 79 | 43.41% | 33 | 18.13% | 58 | 31.87% | 6 | 3.30% | 5 | 2.75% | 1 | 0.55% |
| 40 ～ 74 | 男 | 631 | 121 | 19.18% | 135 | 21.39% | 253 | 40.10% | 42 | 6.66% | 12 | 1.90% | 68 | 10.78% |
| | 女 | 635 | 146 | 22.99% | 126 | 19.84% | 233 | 36.69% | 40 | 6.30% | 15 | 2.36% | 75 | 11.81% |
| | 計 | 1266 | 267 | 21.09% | 261 | 20.62% | 486 | 38.39% | 82 | 6.48% | 27 | 2.13% | 143 | 11.30% |
| 75 ～ | 男 | 22 | 6 | 27.27% | 4 | 18.18% | 8 | 36.36% | 1 | 4.55% | 0 | 0 | 3 | 13.64% |
| | 女 | 12 | 0 | 0.00% | 2 | 16.67% | 3 | 25.00% | 1 | 8.33% | 0 | 0 | 6 | 50.00% |
| | 計 | 34 | 6 | 17.65% | 6 | 17.65% | 11 | 32.35% | 2 | 5.88% | 0 | 0 | 9 | 26.47% |
| 合計 | | 1482 | 352 | 23.75% | 300 | 20.24% | 555 | 37.45% | 90 | 6.07% | 32 | 2.16% | 153 | 10.32% |

6. 糖代謝検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 35 | 40.23% | 39 | 44.83% | 9 | 10.34% | 0 | | 3 | 3.45% | 1 | 1.15% |
| | 女 | 95 | 59 | 62.11% | 33 | 34.74% | 3 | 3.16% | 0 | | 0 | | 0 | |
| | 計 | 182 | 94 | 51.65% | 72 | 39.56% | 12 | 6.59% | 0 | | 3 | 1.65% | 1 | 0.55% |
| 40 ～ 74 | 男 | 631 | 172 | 27.26% | 245 | 38.83% | 145 | 22.98% | 0 | | 35 | 5.55% | 34 | 5.39% |
| | 女 | 635 | 213 | 33.54% | 257 | 40.47% | 132 | 20.79% | 0 | | 17 | 2.68% | 16 | 2.52% |
| | 計 | 1266 | 385 | 30.41% | 502 | 39.65% | 277 | 21.88% | 0 | | 52 | 4.11% | 50 | 3.95% |
| 75 ～ | 男 | 22 | 3 | 13.64% | 10 | 45.45% | 5 | 22.73% | 0 | | 2 | 9.09% | 2 | 9.09% |
| | 女 | 12 | 1 | 8.33% | 2 | 16.67% | 6 | 50.00% | 0 | | 1 | 8.33% | 2 | 16.67% |
| | 計 | 34 | 4 | 11.76% | 12 | 35.29% | 11 | 32.35% | 0 | | 3 | 8.82% | 4 | 11.76% |
| 合計 | | 1482 | 483 | 32.59% | 586 | 39.54% | 300 | 20.24% | 0 | | 58 | 3.91% | 55 | 3.71% |

7. 腎機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 69 | 79.31% | 15 | 17.24% | 3 | 3.45% | 0 | | 0 | | 0 | |
| | 女 | 95 | 57 | 60.00% | 26 | 27.37% | 11 | 11.58% | 0 | | 1 | 1.05% | 0 | |
| | 計 | 182 | 126 | 69.23% | 41 | 22.53% | 14 | 7.69% | 0 | | 1 | 0.55% | 0 | |
| 40 ～ 74 | 男 | 631 | 478 | 75.75% | 88 | 13.95% | 60 | 9.51% | 0 | | 5 | 0.79% | 0 | |
| | 女 | 635 | 377 | 59.37% | 143 | 22.52% | 108 | 17.01% | 0 | | 7 | 1.10% | 0 | |
| | 計 | 1266 | 855 | 67.54% | 231 | 18.25% | 168 | 13.27% | 0 | | 12 | 0.95% | 0 | |
| 75 ～ | 男 | 22 | 12 | 54.55% | 5 | 22.73% | 5 | 22.73% | 0 | | 0 | | 0 | |
| | 女 | 12 | 4 | 33.33% | 2 | 16.67% | 6 | 50.00% | 0 | | 0 | | 0 | |
| | 計 | 34 | 16 | 47.06% | 7 | 20.59% | 11 | 32.35% | 0 | | 0 | | 0 | |
| 合計 | | 1482 | 997 | 67.27% | 279 | 18.83% | 193 | 13.02% | 0 | | 13 | 0.88% | 0 | |

8. 尿酸検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|-------|--------|-----|-------|-------|----|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 64 | 73.56% | 6 | 6.90% | 11 | 12.64% | 5 | 5.75% | 0 | | 1 | 1.15% |
| | 女 | 95 | 94 | 98.95% | 0 | | 1 | 1.05% | 0 | | 0 | | 0 | |
| | 計 | 182 | 158 | 86.81% | 6 | 3.30% | 12 | 6.59% | 5 | 2.75% | 0 | | 1 | 0.55% |
| 40 ～ 74 | 男 | 631 | 490 | 77.65% | 51 | 8.08% | 58 | 9.19% | 4 | 0.63% | 0 | | 28 | 4.44% |
| | 女 | 635 | 623 | 98.11% | 8 | 1.26% | 4 | 0.63% | 0 | | 0 | | 0 | |
| | 計 | 1266 | 1113 | 87.91% | 59 | 4.66% | 62 | 4.90% | 4 | 0.32% | 0 | | 28 | 2.21% |
| 75 ～ | 男 | 22 | 19 | 86.36% | 0 | | 3 | 13.64% | 0 | | 0 | | 0 | |
| | 女 | 12 | 10 | 83.33% | 0 | | 2 | 16.67% | 0 | | 0 | | 0 | |
| | 計 | 34 | 29 | 85.29% | 0 | | 5 | 14.71% | 0 | | 0 | | 0 | |
| 合計 | | 1482 | 1300 | 87.72% | 65 | 4.39% | 79 | 5.33% | 9 | 0.61% | 0 | | 29 | 1.96% |

9. 貧血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 65 | 74.71% | 18 | 20.69% | 0 | | 0 | | 4 | 4.60% | 0 | |
| | 女 | 95 | 69 | 72.63% | 6 | 6.32% | 10 | 10.53% | 8 | 8.42% | 0 | | 2 | 2.11% |
| | 計 | 182 | 134 | 73.63% | 24 | 13.19% | 10 | 5.49% | 8 | 4.40% | 4 | 2.20% | 2 | 1.10% |
| 40 ～ 74 | 男 | 631 | 510 | 80.82% | 84 | 13.31% | 18 | 2.85% | 2 | 0.32% | 17 | 2.69% | 0 | |
| | 女 | 635 | 439 | 69.13% | 86 | 13.54% | 65 | 10.24% | 32 | 5.04% | 4 | 0.63% | 9 | 1.42% |
| | 計 | 1266 | 949 | 74.96% | 170 | 13.43% | 83 | 6.56% | 34 | 2.69% | 21 | 1.66% | 9 | 0.71% |
| 75 ～ | 男 | 22 | 17 | 77.27% | 1 | 4.55% | 3 | 13.64% | 1 | 4.55% | 0 | | 0 | |
| | 女 | 12 | 11 | 91.67% | 1 | 8.33% | 0 | | 0 | | 0 | | 0 | |
| | 計 | 34 | 28 | 82.35% | 2 | 5.88% | 3 | 8.82% | 1 | 2.94% | 0 | | 0 | |
| 合計 | | 1482 | 1111 | 74.97% | 196 | 13.23% | 96 | 6.48% | 43 | 2.90% | 25 | 1.69% | 11 | 0.74% |

10. 血液一般検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 82 | 94.25% | 3 | 3.45% | 0 | | 0 | | 2 | 2.30% | 0 | |
| | 女 | 95 | 79 | 83.16% | 4 | 4.21% | 6 | 6.32% | 0 | | 6 | 6.32% | 0 | |
| | 計 | 182 | 161 | 88.46% | 7 | 3.85% | 6 | 3.30% | 0 | | 8 | 4.40% | 0 | |
| 40 ～ 74 | 男 | 631 | 573 | 90.81% | 18 | 2.85% | 12 | 1.90% | 0 | | 28 | 4.44% | 0 | |
| | 女 | 635 | 562 | 88.50% | 29 | 4.57% | 26 | 4.09% | 0 | | 18 | 2.83% | 0 | |
| | 計 | 1266 | 1135 | 89.65% | 47 | 3.71% | 38 | 3.00% | 0 | | 46 | 3.63% | 0 | |
| 75 ～ | 男 | 22 | 21 | 95.45% | 0 | | 1 | 4.55% | 0 | | 0 | | 0 | |
| | 女 | 12 | 12 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 34 | 33 | 97.06% | 0 | | 1 | 2.94% | 0 | | 0 | | 0 | |
| 合計 | | 1482 | 1329 | 89.68% | 54 | 3.64% | 45 | 3.04% | 0 | | 54 | 3.64% | 0 | |

11. 肝炎検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 5 | 5 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 7 | 7 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 12 | 12 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 41 | 40 | 97.56% | 0 | | 0 | | 0 | | 1 | 2.44% | 0 | |
| | 女 | 13 | 13 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 54 | 53 | 98.15% | 0 | | 0 | | 0 | | 1 | 1.85% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 66 | 65 | 98.48% | 0 | | 0 | | 0 | | 1 | 1.52% | 0 | |

12. 前立腺がん検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 16 | 16 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 16 | 16 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 174 | 165 | 94.83% | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 5.17% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 174 | 165 | 94.83% | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 5.17% | 0 | 0 |
| 75 ～ | 男 | 2 | 2 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 2 | 2 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 192 | 183 | 95.31% | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 4.69% | 0 | 0 |

13. 便潜血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 85 | 83 | 97.65% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2.35% | 0 | 0 |
| | 女 | 93 | 90 | 96.77% | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 3.23% | 0 | 0 |
| | 計 | 178 | 173 | 97.19% | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 2.81% | 0 | 0 |
| 40 ～ 74 | 男 | 621 | 586 | 94.36% | 0 | 0 | 0 | 0 | 0 | 0 | 34 | 5.48% | 1 | 0.16% |
| | 女 | 624 | 600 | 96.15% | 0 | 0 | 0 | 0 | 0 | 0 | 24 | 3.85% | 0 | 0 |
| | 計 | 1245 | 1186 | 95.26% | 0 | 0 | 0 | 0 | 0 | 0 | 58 | 4.66% | 1 | 0.08% |
| 75 ～ | 男 | 22 | 22 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 12 | 12 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 34 | 34 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 1457 | 1393 | 95.61% | 0 | 0 | 0 | 0 | 0 | 0 | 63 | 4.32% | 1 | 0.07% |

14. 心電図検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 74 | 85.06% | 9 | 10.34% | 1 | 1.15% | 0 | 0 | 2 | 2.30% | 1 | 1.15% |
| | 女 | 95 | 85 | 89.47% | 1 | 1.05% | 9 | 9.47% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 182 | 159 | 87.36% | 10 | 5.49% | 10 | 5.49% | 0 | 0 | 2 | 1.10% | 1 | 0.55% |
| 40 ～ 74 | 男 | 629 | 465 | 73.93% | 69 | 10.97% | 63 | 10.02% | 0 | 0 | 16 | 2.54% | 16 | 2.54% |
| | 女 | 635 | 515 | 81.10% | 41 | 6.46% | 67 | 10.55% | 0 | 0 | 5 | 0.79% | 7 | 1.10% |
| | 計 | 1264 | 980 | 77.53% | 110 | 8.70% | 130 | 10.28% | 0 | 0 | 21 | 1.66% | 23 | 1.82% |
| 75 ～ | 男 | 22 | 11 | 50.00% | 4 | 18.18% | 5 | 22.73% | 0 | 0 | 1 | 4.55% | 1 | 4.55% |
| | 女 | 12 | 8 | 66.67% | 2 | 16.67% | 1 | 8.33% | 0 | 0 | 1 | 8.33% | 0 | 0 |
| | 計 | 34 | 19 | 55.88% | 6 | 17.65% | 6 | 17.65% | 0 | 0 | 2 | 5.88% | 1 | 2.94% |
| 合計 | | 1480 | 1158 | 78.24% | 126 | 8.51% | 146 | 9.86% | 0 | 0 | 25 | 1.69% | 25 | 1.69% |

15. 眼底検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 85 | 97.70% | 0 | | 0 | | 0 | | 1 | 1.15% | 1 | 1.15% |
| | 女 | 95 | 94 | 98.95% | 0 | | 0 | | 0 | | 0 | | 1 | 1.05% |
| | 計 | 182 | 179 | 98.35% | 0 | | 0 | | 0 | | 1 | 0.55% | 2 | 1.10% |
| 40 ～ 74 | 男 | 627 | 512 | 81.66% | 24 | 3.83% | 31 | 4.94% | 0 | | 32 | 5.10% | 28 | 4.47% |
| | 女 | 632 | 526 | 83.23% | 29 | 4.59% | 21 | 3.32% | 0 | | 22 | 3.48% | 34 | 5.38% |
| | 計 | 1259 | 1038 | 82.45% | 53 | 4.21% | 52 | 4.13% | 0 | | 54 | 4.29% | 62 | 4.92% |
| 75 ～ | 男 | 22 | 12 | 54.55% | 1 | 4.55% | 5 | 22.73% | 0 | | 1 | 4.55% | 3 | 13.64% |
| | 女 | 12 | 3 | 25.00% | 2 | 16.67% | 3 | 25.00% | 0 | | 1 | 8.33% | 3 | 25.00% |
| | 計 | 34 | 15 | 44.12% | 3 | 8.82% | 8 | 23.53% | 0 | | 2 | 5.88% | 6 | 17.65% |
| 合計 | | 1475 | 1232 | 83.53% | 56 | 3.80% | 60 | 4.07% | 0 | | 57 | 3.86% | 70 | 4.75% |

16. 眼圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|-------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 76 | 87.36% | 0 | | 6 | 6.90% | 0 | | 4 | 4.60% | 1 | 1.15% |
| | 女 | 95 | 86 | 90.53% | 0 | | 3 | 3.16% | 0 | | 5 | 5.26% | 1 | 1.05% |
| | 計 | 182 | 162 | 89.01% | 0 | | 9 | 4.95% | 0 | | 9 | 4.95% | 2 | 1.10% |
| 40 ～ 74 | 男 | 627 | 531 | 84.69% | 0 | | 34 | 5.42% | 0 | | 33 | 5.26% | 29 | 4.63% |
| | 女 | 630 | 553 | 87.78% | 0 | | 22 | 3.49% | 0 | | 21 | 3.33% | 34 | 5.40% |
| | 計 | 1257 | 1084 | 86.24% | 0 | | 56 | 4.46% | 0 | | 54 | 4.30% | 63 | 5.01% |
| 75 ～ | 男 | 22 | 17 | 77.27% | 0 | | 2 | 9.09% | 0 | | 0 | | 3 | 13.64% |
| | 女 | 12 | 9 | 75.00% | 0 | | 0 | | 0 | | 0 | | 3 | 25.00% |
| | 計 | 34 | 26 | 76.47% | 0 | | 2 | 5.88% | 0 | | 0 | | 6 | 17.65% |
| 合計 | | 1473 | 1272 | 86.35% | 0 | | 67 | 4.55% | 0 | | 63 | 4.28% | 71 | 4.82% |

17. 肺機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 82 | 94.25% | 0 | | 0 | | 0 | | 5 | 5.75% | 0 | |
| | 女 | 95 | 90 | 94.74% | 0 | | 0 | | 0 | | 3 | 3.16% | 2 | 2.11% |
| | 計 | 182 | 172 | 94.51% | 0 | | 0 | | 0 | | 8 | 4.40% | 2 | 1.10% |
| 40 ～ 74 | 男 | 631 | 557 | 88.27% | 0 | | 25 | 3.96% | 0 | | 39 | 6.18% | 10 | 1.58% |
| | 女 | 635 | 590 | 92.91% | 0 | | 7 | 1.10% | 0 | | 32 | 5.04% | 6 | 0.94% |
| | 計 | 1266 | 1147 | 90.60% | 0 | | 32 | 2.53% | 0 | | 71 | 5.61% | 16 | 1.26% |
| 75 ～ | 男 | 22 | 12 | 54.55% | 0 | | 5 | 22.73% | 0 | | 5 | 22.73% | 0 | |
| | 女 | 12 | 10 | 83.33% | 0 | | 1 | 8.33% | 0 | | 0 | | 1 | 8.33% |
| | 計 | 34 | 22 | 64.71% | 0 | | 6 | 17.65% | 0 | | 5 | 14.71% | 1 | 2.94% |
| 合計 | | 1482 | 1341 | 90.49% | 0 | | 38 | 2.56% | 0 | | 84 | 5.67% | 19 | 1.28% |

18. 胸部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 78 | 89.66% | 0 | | 9 | 10.34% | 0 | | 0 | | 0 | |
| | 女 | 93 | 82 | 88.17% | 0 | | 10 | 10.75% | 0 | | 1 | 1.08% | 0 | |
| | 計 | 180 | 160 | 88.89% | 0 | | 19 | 10.56% | 0 | | 1 | 0.56% | 0 | |
| 40 ～ 74 | 男 | 629 | 498 | 79.17% | 1 | 0.16% | 122 | 19.40% | 0 | | 4 | 0.64% | 4 | 0.64% |
| | 女 | 630 | 494 | 78.41% | 1 | 0.16% | 132 | 20.95% | 0 | | 2 | 0.32% | 1 | 0.16% |
| | 計 | 1259 | 992 | 78.79% | 2 | 0.16% | 254 | 20.17% | 0 | | 6 | 0.48% | 5 | 0.40% |
| 75 ～ | 男 | 22 | 11 | 50.00% | 0 | | 11 | 50.00% | 0 | | 0 | | 0 | |
| | 女 | 12 | 3 | 25.00% | 0 | | 8 | 66.67% | 0 | | 0 | | 1 | 8.33% |
| | 計 | 34 | 14 | 41.18% | 0 | | 19 | 55.88% | 0 | | 0 | | 1 | 2.94% |
| 合計 | | 1473 | 1166 | 79.16% | 2 | 0.14% | 292 | 19.82% | 0 | | 7 | 0.48% | 6 | 0.41% |

19. 胃部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 73 | 67 | 91.78% | 0 | | 3 | 4.11% | 0 | | 3 | 4.11% | 0 | |
| | 女 | 58 | 49 | 84.48% | 0 | | 8 | 13.79% | 0 | | 1 | 1.72% | 0 | |
| | 計 | 131 | 116 | 88.55% | 0 | | 11 | 8.40% | 0 | | 4 | 3.05% | 0 | |
| 40 ～ 74 | 男 | 357 | 310 | 86.83% | 0 | | 29 | 8.12% | 0 | | 18 | 5.04% | 0 | |
| | 女 | 380 | 315 | 82.89% | 0 | | 60 | 15.79% | 0 | | 4 | 1.05% | 1 | 0.26% |
| | 計 | 737 | 625 | 84.80% | 0 | | 89 | 12.08% | 0 | | 22 | 2.99% | 1 | 0.14% |
| 75 ～ | 男 | 12 | 8 | 66.67% | 0 | | 3 | 25.00% | 0 | | 1 | 8.33% | 0 | |
| | 女 | 6 | 4 | 66.67% | 0 | | 1 | 16.67% | 0 | | 1 | 16.67% | 0 | |
| | 計 | 18 | 12 | 66.67% | 0 | | 4 | 22.22% | 0 | | 2 | 11.11% | 0 | |
| 合計 | | 886 | 753 | 84.99% | 0 | | 104 | 11.74% | 0 | | 28 | 3.16% | 1 | 0.11% |

20. 胃内視鏡検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|----|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 11 | 6 | 54.55% | 0 | | 5 | 45.45% | 0 | | 0 | | 0 | |
| | 女 | 27 | 13 | 48.15% | 0 | | 14 | 51.85% | 0 | | 0 | | 0 | |
| | 計 | 38 | 19 | 50.00% | 0 | | 19 | 50.00% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 251 | 43 | 17.13% | 0 | | 178 | 70.92% | 0 | | 28 | 11.16% | 2 | 0.80% |
| | 女 | 194 | 44 | 22.68% | 1 | 0.52% | 134 | 69.07% | 0 | | 14 | 7.22% | 1 | 0.52% |
| | 計 | 445 | 87 | 19.55% | 1 | 0.22% | 312 | 70.11% | 0 | | 42 | 9.44% | 3 | 0.67% |
| 75 ～ | 男 | 10 | 0 | | 0 | | 8 | 80.00% | 0 | | 2 | 20.00% | 0 | |
| | 女 | 5 | 0 | | 0 | | 3 | 60.00% | 0 | | 2 | 40.00% | 0 | |
| | 計 | 15 | 0 | | 0 | | 11 | 73.33% | 0 | | 4 | 26.67% | 0 | |
| 合計 | | 498 | 106 | 21.29% | 1 | 0.20% | 342 | 68.67% | 0 | | 46 | 9.24% | 3 | 0.60% |

21. 腹部超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 87 | 21 | 24.14% | 1 | 1.15% | 63 | 72.41% | 0 | 2 | 2.30% | 0 | |
| | 女 | 95 | 48 | 50.53% | 6 | 6.32% | 40 | 42.11% | 0 | 1 | 1.05% | 0 | |
| | 計 | 182 | 69 | 37.91% | 7 | 3.85% | 103 | 56.59% | 0 | 3 | 1.65% | 0 | |
| 40 ～ 74 | 男 | 631 | 67 | 10.62% | 1 | 0.16% | 530 | 83.99% | 0 | 31 | 4.91% | 2 | 0.32% |
| | 女 | 635 | 119 | 18.74% | 5 | 0.79% | 481 | 75.75% | 0 | 30 | 4.72% | 0 | |
| | 計 | 1266 | 186 | 14.69% | 6 | 0.47% | 1011 | 79.86% | 0 | 61 | 4.82% | 2 | 0.16% |
| 75 ～ | 男 | 22 | 0 | | 0 | | 19 | 86.36% | 0 | 3 | 13.64% | 0 | |
| | 女 | 12 | 1 | 8.33% | 0 | | 9 | 75.00% | 0 | 2 | 16.67% | 0 | |
| | 計 | 34 | 1 | 2.94% | 0 | | 28 | 82.35% | 0 | 5 | 14.71% | 0 | |
| 合計 | | 1482 | 256 | 17.27% | 13 | 0.88% | 1142 | 77.06% | 0 | 69 | 4.66% | 2 | 0.13% |

22. 乳房マンモグラフィ検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|-------|-----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 28 | 24 | 85.71% | 0 | | 1 | 3.57% | 0 | 3 | 10.71% | 0 | |
| | 計 | 28 | 24 | 85.71% | 0 | | 1 | 3.57% | 0 | 3 | 10.71% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 179 | 158 | 88.27% | 0 | | 15 | 8.38% | 0 | 6 | 3.35% | 0 | |
| | 計 | 179 | 158 | 88.27% | 0 | | 15 | 8.38% | 0 | 6 | 3.35% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| 合計 | | 207 | 182 | 87.92% | 0 | | 16 | 7.73% | 0 | 9 | 4.35% | 0 | |

23. 乳房超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|--------|-----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 48 | 35 | 72.92% | 0 | | 12 | 25.00% | 0 | 1 | 2.08% | 0 | |
| | 計 | 48 | 35 | 72.92% | 0 | | 12 | 25.00% | 0 | 1 | 2.08% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 214 | 114 | 53.27% | 0 | | 92 | 42.99% | 0 | 8 | 3.74% | 0 | |
| | 計 | 214 | 114 | 53.27% | 0 | | 92 | 42.99% | 0 | 8 | 3.74% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | 0 | 0 | | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | 0 | 0 | | |
| 合計 | | 263 | 150 | 57.03% | 0 | | 104 | 39.54% | 0 | 9 | 3.42% | 0 | |

24. 婦人科検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|-------|-----|----|-------|-------|-----|-------|
| | | | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 45 | 43 | 95.56% | 0 | | 0 | | 0 | | 2 | 4.44% | 0 | |
| | 計 | 45 | 43 | 95.56% | 0 | | 0 | | 0 | | 2 | 4.44% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 248 | 239 | 96.37% | 3 | 1.21% | 3 | 1.21% | 0 | | 1 | 0.40% | 2 | 0.81% |
| | 計 | 248 | 239 | 96.37% | 3 | 1.21% | 3 | 1.21% | 0 | | 1 | 0.40% | 2 | 0.81% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 293 | 282 | 96.25% | 3 | 1.02% | 3 | 1.02% | 0 | | 3 | 1.02% | 2 | 0.68% |

②生活習慣病健診

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|-------|------|--------|---------------|--------|-----|--------|-------|--------|------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 63 | 9.13% | 72 | 10.43% | 294 | 42.61% | 33 | 4.78% | 192 | 27.83% | 36 | 5.22% |
| | 女 | 612 | 57 | 9.31% | 56 | 9.15% | 306 | 50.00% | 66 | 10.78% | 108 | 17.65% | 19 | 3.10% |
| | 計 | 1302 | 120 | 9.22% | 128 | 9.83% | 600 | 46.08% | 99 | 7.60% | 300 | 23.04% | 55 | 4.22% |
| 40 ～ 74 | 男 | 2417 | 26 | 1.08% | 668 | 27.64% | 819 | 33.88% | 130 | 5.38% | 672 | 27.80% | 668 | 27.64% |
| | 女 | 2299 | 39 | 1.70% | 424 | 18.44% | 1045 | 45.45% | 218 | 9.48% | 467 | 20.31% | 424 | 18.44% |
| | 計 | 4716 | 65 | 1.38% | 1092 | 23.16% | 1864 | 39.53% | 348 | 7.38% | 1139 | 24.15% | 1092 | 23.16% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 6018 | 185 | 3.07% | 1220 | 20.27% | 2464 | 40.94% | 447 | 7.43% | 1439 | 23.91% | 1147 | 19.06% |

1. BMI

| 年代 | 性別 | 受診者数 | ～ 18.4 | | 18.5 ～ 24.9 | | 25.0 ～ | |
|---------|----|------|--------|--------|-------------|--------|--------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 51 | 7.39% | 402 | 58.26% | 237 | 34.35% |
| | 女 | 612 | 101 | 16.50% | 416 | 67.97% | 95 | 15.52% |
| | 計 | 1302 | 152 | 11.67% | 818 | 62.83% | 332 | 25.50% |
| 40 ～ 74 | 男 | 2417 | 86 | 3.56% | 1496 | 61.89% | 835 | 34.55% |
| | 女 | 2299 | 299 | 13.01% | 1566 | 68.12% | 434 | 18.88% |
| | 計 | 4716 | 385 | 8.16% | 3062 | 64.93% | 1269 | 26.91% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | |
| 合計 | | 6018 | 537 | 8.92% | 3880 | 64.47% | 1601 | 26.60% |

2. 血圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 472 | 68.41% | 114 | 16.52% | 67 | 9.71% | 0 | | 19 | 2.75% | 18 | 2.61% |
| | 女 | 612 | 549 | 89.71% | 29 | 4.74% | 25 | 4.08% | 0 | | 6 | 0.98% | 3 | 0.49% |
| | 計 | 1302 | 1021 | 78.42% | 143 | 10.98% | 92 | 7.07% | 0 | | 25 | 1.92% | 21 | 1.61% |
| 40 ～ 74 | 男 | 2417 | 1027 | 42.49% | 380 | 15.72% | 395 | 16.34% | 0 | | 156 | 6.45% | 459 | 18.99% |
| | 女 | 2299 | 1444 | 62.81% | 277 | 12.05% | 262 | 11.40% | 0 | | 70 | 3.04% | 246 | 10.70% |
| | 計 | 4716 | 2471 | 52.40% | 657 | 13.93% | 657 | 13.93% | 0 | | 226 | 4.79% | 705 | 14.95% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 6018 | 3492 | 58.03% | 800 | 13.29% | 749 | 12.45% | 0 | | 251 | 4.17% | 726 | 12.06% |

3. 聴力検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 689 | 675 | 97.97% | 0 | | 10 | 1.45% | 4 | 0.58% | 0 | | 0 | |
| | 女 | 589 | 579 | 98.30% | 0 | | 3 | 0.51% | 6 | 1.02% | 0 | | 1 | 0.17% |
| | 計 | 1278 | 1254 | 98.12% | 0 | | 13 | 1.02% | 10 | 0.78% | 0 | | 1 | 0.08% |
| 40 ～ 74 | 男 | 2404 | 2097 | 87.23% | 0 | | 265 | 11.02% | 40 | 1.66% | 0 | | 2 | 0.08% |
| | 女 | 2220 | 2122 | 95.59% | 0 | | 79 | 3.56% | 16 | 0.72% | 1 | 0.05% | 2 | 0.09% |
| | 計 | 4624 | 4219 | 91.24% | 0 | | 344 | 7.44% | 56 | 1.21% | 1 | 0.02% | 4 | 0.09% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5902 | 5473 | 92.73% | 0 | | 357 | 6.05% | 66 | 1.12% | 1 | 0.02% | 5 | 0.08% |

4. 肝機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 638 | 360 | 56.43% | 100 | 15.67% | 63 | 9.87% | 0 | | 115 | 18.03% | 0 | |
| | 女 | 576 | 514 | 89.24% | 25 | 4.34% | 17 | 2.95% | 0 | | 20 | 3.47% | 0 | |
| | 計 | 1214 | 874 | 71.99% | 125 | 10.30% | 80 | 6.59% | 0 | | 135 | 11.12% | 0 | |
| 40 ～ 74 | 男 | 2416 | 1289 | 53.35% | 455 | 18.83% | 278 | 11.51% | 0 | | 384 | 15.89% | 10 | 0.41% |
| | 女 | 2299 | 1843 | 80.17% | 198 | 8.61% | 126 | 5.48% | 1 | 0.04% | 128 | 5.57% | 3 | 0.13% |
| | 計 | 4715 | 3132 | 66.43% | 653 | 13.85% | 404 | 8.57% | 1 | 0.02% | 512 | 10.86% | 13 | 0.28% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5929 | 4006 | 67.57% | 778 | 13.12% | 484 | 8.16% | 1 | 0.02% | 647 | 10.91% | 13 | 0.22% |

5. 脂質検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 638 | 244 | 38.24% | 133 | 20.85% | 187 | 29.31% | 49 | 7.68% | 13 | 2.04% | 12 | 1.88% |
| | 女 | 576 | 345 | 59.90% | 98 | 17.01% | 99 | 17.19% | 10 | 1.74% | 24 | 4.17% | 0 | |
| | 計 | 1214 | 589 | 48.52% | 231 | 19.03% | 286 | 23.56% | 59 | 4.86% | 37 | 3.05% | 12 | 0.99% |
| 40 ～ 74 | 男 | 2416 | 577 | 23.88% | 555 | 22.97% | 830 | 34.35% | 216 | 8.94% | 47 | 1.95% | 191 | 7.91% |
| | 女 | 2299 | 725 | 31.54% | 481 | 20.92% | 681 | 29.62% | 163 | 7.09% | 54 | 2.35% | 195 | 8.48% |
| | 計 | 4715 | 1302 | 27.61% | 1036 | 21.97% | 1511 | 32.05% | 379 | 8.04% | 101 | 2.14% | 386 | 8.19% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5929 | 1891 | 31.89% | 1267 | 21.37% | 1797 | 30.31% | 438 | 7.39% | 138 | 2.33% | 398 | 6.71% |

6. 糖代謝検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 487 | 70.58% | 152 | 22.03% | 25 | 3.62% | 0 | 0 | 11 | 1.59% | 15 | 2.17% |
| | 女 | 611 | 482 | 78.89% | 113 | 18.49% | 13 | 2.13% | 0 | 0 | 3 | 0.49% | 0 | 0 |
| | 計 | 1301 | 969 | 74.48% | 265 | 20.37% | 38 | 2.92% | 0 | 0 | 14 | 1.08% | 15 | 1.15% |
| 40 ～ 74 | 男 | 2417 | 1219 | 50.43% | 630 | 26.07% | 276 | 11.42% | 0 | 0 | 131 | 5.42% | 161 | 6.66% |
| | 女 | 2299 | 1390 | 60.46% | 587 | 25.53% | 227 | 9.87% | 0 | 0 | 35 | 1.52% | 60 | 2.61% |
| | 計 | 4716 | 2609 | 55.32% | 1217 | 25.81% | 503 | 10.67% | 0 | 0 | 166 | 3.52% | 221 | 4.69% |
| 75 ～ | 男 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 6017 | 3578 | 59.46% | 1482 | 24.63% | 541 | 8.99% | 0 | 0 | 180 | 2.99% | 236 | 3.92% |

7. 腎機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 594 | 86.09% | 60 | 8.70% | 29 | 4.20% | 0 | 0 | 7 | 1.01% | 0 | 0 |
| | 女 | 611 | 431 | 70.54% | 116 | 18.99% | 60 | 9.82% | 0 | 0 | 4 | 0.65% | 0 | 0 |
| | 計 | 1301 | 1025 | 78.79% | 176 | 13.53% | 89 | 6.84% | 0 | 0 | 11 | 0.85% | 0 | 0 |
| 40 ～ 74 | 男 | 2417 | 1953 | 80.80% | 239 | 9.89% | 180 | 7.45% | 0 | 0 | 45 | 1.86% | 0 | 0 |
| | 女 | 2299 | 1469 | 63.90% | 503 | 21.88% | 306 | 13.31% | 0 | 0 | 21 | 0.91% | 0 | 0 |
| | 計 | 4716 | 3422 | 72.56% | 742 | 15.73% | 486 | 10.31% | 0 | 0 | 66 | 1.40% | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 6017 | 4447 | 73.91% | 918 | 15.26% | 575 | 9.56% | 0 | 0 | 77 | 1.28% | 0 | 0 |

8. 尿酸検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 594 | 454 | 76.43% | 59 | 9.93% | 65 | 10.94% | 9 | 1.52% | 5 | 0.84% | 2 | 0.34% |
| | 女 | 529 | 525 | 99.24% | 2 | 0.38% | 2 | 0.38% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 1123 | 979 | 87.18% | 61 | 5.43% | 67 | 5.97% | 9 | 0.80% | 5 | 0.45% | 2 | 0.18% |
| 40 ～ 74 | 男 | 2404 | 1876 | 78.04% | 219 | 9.11% | 235 | 9.78% | 31 | 1.29% | 8 | 0.33% | 35 | 1.46% |
| | 女 | 2250 | 2204 | 97.96% | 20 | 0.89% | 24 | 1.07% | 0 | 0 | 1 | 0.04% | 1 | 0.04% |
| | 計 | 4654 | 4080 | 87.67% | 239 | 5.14% | 259 | 5.57% | 31 | 0.67% | 9 | 0.19% | 36 | 0.77% |
| 75 ～ | 男 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 5777 | 5059 | 87.57% | 300 | 5.19% | 326 | 5.64% | 40 | 0.69% | 14 | 0.24% | 38 | 0.66% |

9. 貧血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|--------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 623 | 467 | 74.96% | 113 | 18.14% | 9 | 1.44% | 5 | 0.80% | 29 | 4.65% | 0 | |
| | 女 | 597 | 370 | 61.98% | 54 | 9.05% | 96 | 16.08% | 61 | 10.22% | 7 | 1.17% | 9 | 1.51% |
| | 計 | 1220 | 837 | 68.61% | 167 | 13.69% | 105 | 8.61% | 66 | 5.41% | 36 | 2.95% | 9 | 0.74% |
| 40 ～ 74 | 男 | 2416 | 1881 | 77.86% | 304 | 12.58% | 93 | 3.85% | 28 | 1.16% | 105 | 4.35% | 5 | 0.21% |
| | 女 | 2299 | 1580 | 68.73% | 250 | 10.87% | 276 | 12.01% | 137 | 5.96% | 20 | 0.87% | 36 | 1.57% |
| | 計 | 4715 | 3461 | 73.40% | 554 | 11.75% | 369 | 7.83% | 165 | 3.50% | 125 | 2.65% | 41 | 0.87% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5935 | 4298 | 72.42% | 721 | 12.15% | 474 | 7.99% | 231 | 3.89% | 161 | 2.71% | 50 | 0.84% |

10. 血液一般検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 456 | 409 | 89.69% | 10 | 2.19% | 2 | 0.44% | 0 | | 35 | 7.68% | 0 | |
| | 女 | 392 | 357 | 91.07% | 16 | 4.08% | 4 | 1.02% | 0 | | 15 | 3.83% | 0 | |
| | 計 | 848 | 766 | 90.33% | 26 | 3.07% | 6 | 0.71% | 0 | | 50 | 5.90% | 0 | |
| 40 ～ 74 | 男 | 1952 | 1740 | 89.14% | 60 | 3.07% | 24 | 1.23% | 0 | | 128 | 6.56% | 0 | |
| | 女 | 1894 | 1728 | 91.24% | 49 | 2.59% | 60 | 3.17% | 0 | | 57 | 3.01% | 0 | |
| | 計 | 3846 | 3468 | 90.17% | 109 | 2.83% | 84 | 2.18% | 0 | | 185 | 4.81% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 4694 | 4234 | 90.20% | 135 | 2.88% | 90 | 1.92% | 0 | | 235 | 5.01% | 0 | |

11. 肝炎検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 50 | 50 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 42 | 41 | 97.62% | 0 | | 0 | | 0 | | 1 | 2.38% | 0 | |
| | 計 | 92 | 91 | 98.91% | 0 | | 0 | | 0 | | 1 | 1.09% | 0 | |
| 40 ～ 74 | 男 | 360 | 358 | 99.44% | 0 | | 0 | | 0 | | 2 | 0.56% | 0 | |
| | 女 | 284 | 279 | 98.24% | 0 | | 0 | | 0 | | 5 | 1.76% | 0 | |
| | 計 | 644 | 637 | 98.91% | 0 | | 0 | | 0 | | 7 | 1.09% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 736 | 728 | 98.91% | 0 | | 0 | | 0 | | 8 | 1.09% | 0 | |

12. 前立腺がん検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 11 | 11 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 11 | 11 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 274 | 266 | 97.08% | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 2.92% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 274 | 266 | 97.08% | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 2.92% | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 285 | 277 | 97.19% | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 2.81% | 0 | 0 |

13. 便潜血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|----|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 537 | 514 | 95.72% | 0 | 0 | 0 | 0 | 0 | 0 | 23 | 4.28% | 0 | 0 |
| | 女 | 433 | 416 | 96.07% | 0 | 0 | 0 | 0 | 0 | 0 | 14 | 3.23% | 3 | 0.69% |
| | 計 | 970 | 930 | 95.88% | 0 | 0 | 0 | 0 | 0 | 0 | 37 | 3.81% | 3 | 0.31% |
| 40 ～ 74 | 男 | 2313 | 2201 | 95.16% | 0 | 0 | 0 | 0 | 0 | 0 | 109 | 4.71% | 3 | 0.13% |
| | 女 | 2135 | 2050 | 96.02% | 0 | 0 | 0 | 0 | 0 | 0 | 84 | 3.93% | 1 | 0.05% |
| | 計 | 4448 | 4251 | 95.57% | 0 | 0 | 0 | 0 | 0 | 0 | 193 | 4.34% | 4 | 0.09% |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 5418 | 5181 | 95.63% | 0 | 0 | 0 | 0 | 0 | 0 | 230 | 4.25% | 7 | 0.13% |

14. 心電図検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|-------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 607 | 517 | 85.17% | 55 | 9.06% | 29 | 4.78% | 0 | 0 | 5 | 0.82% | 1 | 0.16% |
| | 女 | 536 | 467 | 87.13% | 26 | 4.85% | 37 | 6.90% | 0 | 0 | 6 | 1.12% | 0 | 0 |
| | 計 | 1143 | 984 | 86.09% | 81 | 7.09% | 66 | 5.77% | 0 | 0 | 11 | 0.96% | 1 | 0.09% |
| 40 ～ 74 | 男 | 2416 | 1871 | 77.44% | 214 | 8.86% | 216 | 8.94% | 6 | 0.25% | 71 | 2.94% | 38 | 1.57% |
| | 女 | 2298 | 1957 | 85.16% | 141 | 6.14% | 150 | 6.53% | 1 | 0.04% | 33 | 1.44% | 16 | 0.70% |
| | 計 | 4714 | 3828 | 81.20% | 355 | 7.53% | 366 | 7.76% | 7 | 0.15% | 104 | 2.21% | 54 | 1.15% |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 5857 | 4812 | 82.16% | 436 | 7.44% | 432 | 7.38% | 7 | 0.12% | 115 | 1.96% | 55 | 0.94% |

15. 眼底検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|-------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 141 | 135 | 95.74% | 1 | 0.71% | 0 | | 2 | 1.42% | 3 | 2.13% | 0 | |
| | 女 | 82 | 80 | 97.56% | 1 | 1.22% | 0 | | 0 | | 0 | | 1 | 1.22% |
| | 計 | 223 | 215 | 96.41% | 2 | 0.90% | 0 | | 2 | 0.90% | 3 | 1.35% | 1 | 0.45% |
| 40 ～ 74 | 男 | 555 | 469 | 84.50% | 17 | 3.06% | 27 | 4.86% | 13 | 2.34% | 14 | 2.52% | 15 | 2.70% |
| | 女 | 466 | 405 | 86.91% | 13 | 2.79% | 17 | 3.65% | 5 | 1.07% | 17 | 3.65% | 9 | 1.93% |
| | 計 | 1021 | 874 | 85.60% | 30 | 2.94% | 44 | 4.31% | 18 | 1.76% | 31 | 3.04% | 24 | 2.35% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1244 | 1089 | 87.54% | 32 | 2.57% | 44 | 3.54% | 20 | 1.61% | 34 | 2.73% | 25 | 2.01% |

16. 眼圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 76 | 72 | 94.74% | 0 | | 3 | 3.95% | 0 | | 1 | 1.32% | 0 | |
| | 女 | 39 | 33 | 84.62% | 0 | | 4 | 10.26% | 0 | | 1 | 2.56% | 1 | 2.56% |
| | 計 | 115 | 105 | 91.30% | 0 | | 7 | 6.09% | 0 | | 2 | 1.74% | 1 | 0.87% |
| 40 ～ 74 | 男 | 299 | 259 | 86.62% | 0 | | 16 | 5.35% | 9 | 3.01% | 7 | 2.34% | 8 | 2.68% |
| | 女 | 233 | 213 | 91.42% | 0 | | 7 | 3.00% | 5 | 2.15% | 5 | 2.15% | 3 | 1.29% |
| | 計 | 532 | 472 | 88.72% | 0 | | 23 | 4.32% | 14 | 2.63% | 12 | 2.26% | 11 | 2.07% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 647 | 577 | 89.18% | 0 | | 30 | 4.64% | 14 | 2.16% | 14 | 2.16% | 12 | 1.85% |

17. 肺機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|-------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 110 | 106 | 96.36% | 0 | | 2 | 1.82% | 0 | | 2 | 1.82% | 0 | |
| | 女 | 53 | 52 | 98.11% | 0 | | 0 | | 0 | | 1 | 1.89% | 0 | |
| | 計 | 163 | 158 | 96.93% | 0 | | 2 | 1.23% | 0 | | 3 | 1.84% | 0 | |
| 40 ～ 74 | 男 | 390 | 353 | 90.51% | 0 | | 13 | 3.33% | 0 | | 17 | 4.36% | 7 | 1.79% |
| | 女 | 246 | 228 | 92.68% | 0 | | 4 | 1.63% | 0 | | 12 | 4.88% | 2 | 0.81% |
| | 計 | 636 | 581 | 91.35% | 0 | | 17 | 2.67% | 0 | | 29 | 4.56% | 9 | 1.42% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 799 | 739 | 92.49% | 0 | | 19 | 2.38% | 0 | | 32 | 4.01% | 9 | 1.13% |

18. 胸部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 690 | 604 | 87.54% | 0 | | 85 | 12.32% | 0 | | 1 | 0.14% | 0 | |
| | 女 | 590 | 532 | 90.17% | 0 | | 57 | 9.66% | 0 | | 1 | 0.17% | 0 | |
| | 計 | 1280 | 1136 | 88.75% | 0 | | 142 | 11.09% | 0 | | 2 | 0.16% | 0 | |
| 40 ～ 74 | 男 | 2413 | 1930 | 79.98% | 1 | 0.04% | 466 | 19.31% | 0 | | 12 | 0.50% | 4 | 0.17% |
| | 女 | 2256 | 1886 | 83.60% | 0 | | 358 | 15.87% | 0 | | 12 | 0.53% | 0 | |
| | 計 | 4669 | 3816 | 81.73% | 1 | 0.02% | 824 | 17.65% | 0 | | 24 | 0.51% | 4 | 0.09% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5949 | 4952 | 83.24% | 1 | 0.02% | 966 | 16.24% | 0 | | 26 | 0.44% | 4 | 0.07% |

19. 胃部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 474 | 447 | 94.30% | 0 | | 20 | 4.22% | 0 | | 7 | 1.48% | 0 | |
| | 女 | 322 | 280 | 86.96% | 0 | | 39 | 12.11% | 0 | | 3 | 0.93% | 0 | |
| | 計 | 796 | 727 | 91.33% | 0 | | 59 | 7.41% | 0 | | 10 | 1.26% | 0 | |
| 40 ～ 74 | 男 | 1893 | 1647 | 87.00% | 0 | | 158 | 8.35% | 0 | | 79 | 4.17% | 9 | 0.48% |
| | 女 | 1499 | 1230 | 82.05% | 0 | | 241 | 16.08% | 0 | | 26 | 1.73% | 2 | 0.13% |
| | 計 | 3392 | 2877 | 84.82% | 0 | | 399 | 11.76% | 0 | | 105 | 3.10% | 11 | 0.32% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 4188 | 3604 | 86.06% | 0 | | 458 | 10.94% | 0 | | 115 | 2.75% | 11 | 0.26% |

20. 胃内視鏡検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|-------|-------|--------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 47 | 16 | 34.04% | 0 | | 25 | 53.19% | 0 | | 6 | 12.77% | 0 | |
| | 女 | 44 | 17 | 38.64% | 0 | | 26 | 59.09% | 0 | | 1 | 2.27% | 0 | |
| | 計 | 91 | 33 | 36.26% | 0 | | 51 | 56.04% | 0 | | 7 | 7.69% | 0 | |
| 40 ～ 74 | 男 | 276 | 58 | 21.01% | 1 | 0.36% | 191 | 69.20% | 1 | 0.36% | 24 | 8.70% | 1 | 0.36% |
| | 女 | 195 | 60 | 30.77% | 1 | 0.51% | 117 | 60.00% | 1 | 0.51% | 16 | 8.21% | 0 | |
| | 計 | 471 | 118 | 25.05% | 2 | 0.42% | 308 | 65.39% | 2 | 0.42% | 40 | 8.49% | 1 | 0.21% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 562 | 151 | 26.87% | 2 | 0.36% | 359 | 63.88% | 2 | 0.36% | 47 | 8.36% | 1 | 0.18% |

21. 腹部超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 121 | 26 | 21.49% | 4 | 3.31% | 86 | 71.07% | 0 | | 4 | 3.31% | 1 | 0.83% |
| | 女 | 79 | 25 | 31.65% | 2 | 2.53% | 49 | 62.03% | 1 | 1.27% | 2 | 2.53% | 0 | |
| | 計 | 200 | 51 | 25.50% | 6 | 3.00% | 135 | 67.50% | 1 | 0.50% | 6 | 3.00% | 1 | 0.50% |
| 40 ～ 74 | 男 | 496 | 56 | 11.29% | 9 | 1.81% | 398 | 80.24% | 13 | 2.62% | 20 | 4.03% | 0 | |
| | 女 | 432 | 84 | 19.44% | 9 | 2.08% | 319 | 73.84% | 9 | 2.08% | 11 | 2.55% | 0 | |
| | 計 | 928 | 140 | 15.09% | 18 | 1.94% | 717 | 77.26% | 22 | 2.37% | 31 | 3.34% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1128 | 191 | 16.93% | 24 | 2.13% | 852 | 75.53% | 23 | 2.04% | 37 | 3.28% | 1 | 0.09% |

22. 乳房マンモグラフィ検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 99 | 93 | 93.94% | 0 | | 2 | 2.02% | 0 | | 4 | 4.04% | 0 | |
| | 計 | 99 | 93 | 93.94% | 0 | | 2 | 2.02% | 0 | | 4 | 4.04% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 682 | 588 | 86.22% | 0 | | 47 | 6.89% | 0 | | 47 | 6.89% | 0 | |
| | 計 | 682 | 588 | 86.22% | 0 | | 47 | 6.89% | 0 | | 47 | 6.89% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 781 | 681 | 87.20% | 0 | | 49 | 6.27% | 0 | | 51 | 6.53% | 0 | |

23. 乳房超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 122 | 74 | 60.66% | 0 | | 46 | 37.70% | 0 | | 2 | 1.64% | 0 | |
| | 計 | 122 | 74 | 60.66% | 0 | | 46 | 37.70% | 0 | | 2 | 1.64% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 461 | 249 | 54.01% | 0 | | 198 | 42.95% | 0 | | 14 | 3.04% | 0 | |
| | 計 | 461 | 249 | 54.01% | 0 | | 198 | 42.95% | 0 | | 14 | 3.04% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 583 | 323 | 55.40% | 0 | | 244 | 41.85% | 0 | | 16 | 2.74% | 0 | |

24. 婦人科検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|-------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 202 | 191 | 94.55% | 2 | 0.99% | 1 | 0.50% | 0 | | 6 | 2.97% | 2 | 0.99% |
| | 計 | 202 | 191 | 94.55% | 2 | 0.99% | 1 | 0.50% | 0 | | 6 | 2.97% | 2 | 0.99% |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 853 | 825 | 96.72% | 16 | 1.88% | 0 | | 0 | | 10 | 1.17% | 2 | 0.23% |
| | 計 | 853 | 825 | 96.72% | 16 | 1.88% | 0 | | 0 | | 10 | 1.17% | 2 | 0.23% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1055 | 1016 | 96.30% | 18 | 1.71% | 1 | 0.09% | 0 | | 16 | 1.52% | 4 | 0.38% |

③特定健診

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|--------|------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 1 | 33.33% | 0 | | 1 | 33.33% | 0 | | 0 | | 1 | 33.33% |
| | 女 | 31 | 6 | 19.35% | 3 | 9.68% | 13 | 41.94% | 3 | 9.68% | 4 | 12.90% | 2 | 6.45% |
| | 計 | 34 | 7 | 20.59% | 3 | 8.82% | 14 | 41.18% | 3 | 8.82% | 4 | 11.76% | 3 | 8.82% |
| 40 ～ 74 | 男 | 804 | 4 | 0.50% | 383 | 47.64% | 225 | 27.99% | 23 | 2.86% | 128 | 15.92% | 383 | 47.64% |
| | 女 | 2123 | 109 | 5.13% | 608 | 28.64% | 821 | 38.67% | 127 | 5.98% | 277 | 13.05% | 608 | 28.64% |
| | 計 | 2927 | 113 | 3.86% | 991 | 33.86% | 1046 | 35.74% | 150 | 5.12% | 405 | 13.84% | 991 | 33.86% |
| 75 ～ | 男 | 242 | 0 | | 143 | 59.09% | 52 | 21.49% | 5 | 2.07% | 40 | 16.53% | 143 | 59.09% |
| | 女 | 249 | 0 | | 164 | 65.86% | 43 | 17.27% | 3 | 1.20% | 31 | 12.45% | 164 | 65.86% |
| | 計 | 491 | 0 | | 307 | 62.53% | 95 | 19.35% | 8 | 1.63% | 71 | 14.46% | 307 | 62.53% |
| 合計 | | 3452 | 120 | 3.48% | 1301 | 37.69% | 1155 | 33.46% | 161 | 4.66% | 480 | 13.90% | 1301 | 37.69% |

1. BMI

| 年代 | 性別 | 受診者数 | ～ 18.4 | | 18.5 ～ 24.9 | | 25.0 ～ | |
|---------|----|------|--------|--------|-------------|--------|--------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 0 | | 2 | 66.67% | 1 | 33.33% |
| | 女 | 31 | 5 | 16.13% | 19 | 61.29% | 7 | 22.58% |
| | 計 | 34 | 5 | 14.71% | 21 | 61.76% | 8 | 23.53% |
| 40 ～ 74 | 男 | 804 | 25 | 3.11% | 591 | 73.51% | 188 | 23.38% |
| | 女 | 2123 | 232 | 10.93% | 1488 | 70.09% | 403 | 18.98% |
| | 計 | 2927 | 257 | 8.78% | 2079 | 71.03% | 591 | 20.19% |
| 75 ～ | 男 | 242 | 7 | 2.89% | 202 | 83.47% | 33 | 13.64% |
| | 女 | 249 | 27 | 10.84% | 175 | 70.28% | 47 | 18.88% |
| | 計 | 491 | 34 | 6.92% | 377 | 76.78% | 80 | 16.29% |
| 合計 | | 3452 | 296 | 8.57% | 2477 | 71.76% | 679 | 19.67% |

2. 血圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 2 | 66.67% | 1 | 33.33% | 0 | | 0 | | 0 | | 0 | |
| | 女 | 31 | 25 | 80.65% | 1 | 3.23% | 4 | 12.90% | 0 | | 0 | | 1 | 3.23% |
| | 計 | 34 | 27 | 79.41% | 2 | 5.88% | 4 | 11.76% | 0 | | 0 | | 1 | 2.94% |
| 40 ～ 74 | 男 | 804 | 229 | 28.48% | 113 | 14.05% | 137 | 17.04% | 0 | | 46 | 5.72% | 279 | 34.70% |
| | 女 | 2123 | 1056 | 49.74% | 265 | 12.48% | 282 | 13.28% | 0 | | 117 | 5.51% | 403 | 18.98% |
| | 計 | 2927 | 1285 | 43.90% | 378 | 12.91% | 419 | 14.31% | 0 | | 163 | 5.57% | 682 | 23.30% |
| 75 ～ | 男 | 242 | 38 | 15.70% | 20 | 8.26% | 50 | 20.66% | 0 | | 21 | 8.68% | 113 | 46.69% |
| | 女 | 249 | 32 | 12.85% | 28 | 11.24% | 52 | 20.88% | 0 | | 14 | 5.62% | 123 | 49.40% |
| | 計 | 491 | 70 | 14.26% | 48 | 9.78% | 102 | 20.77% | 0 | | 35 | 7.13% | 236 | 48.07% |
| 合計 | | 3452 | 1382 | 40.03% | 428 | 12.40% | 525 | 15.21% | 0 | | 198 | 5.74% | 919 | 26.62% |

3. 聴力検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|--------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 3 | 1 | 33.33% | 0 | | 2 | 66.67% | 0 | | 0 | | 0 | |
| | 女 | 17 | 15 | 88.24% | 0 | | 2 | 11.76% | 0 | | 0 | | 0 | |
| | 計 | 20 | 16 | 80.00% | 0 | | 4 | 20.00% | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 3 | 3 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 23 | 19 | 82.61% | 0 | | 4 | 17.39% | 0 | | 0 | | 0 | |

4. 肝機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 1 | 33.33% | 1 | 33.33% | 1 | 33.33% | 0 | | 0 | | 0 | |
| | 女 | 31 | 27 | 87.10% | 4 | 12.90% | 0 | | 0 | | 0 | | 0 | |
| | 計 | 34 | 28 | 82.35% | 5 | 14.71% | 1 | 2.94% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 804 | 532 | 66.17% | 131 | 16.29% | 61 | 7.59% | 0 | | 79 | 9.83% | 1 | 0.12% |
| | 女 | 2123 | 1826 | 86.01% | 154 | 7.25% | 69 | 3.25% | 0 | | 71 | 3.34% | 3 | 0.14% |
| | 計 | 2927 | 2358 | 80.56% | 285 | 9.74% | 130 | 4.44% | 0 | | 150 | 5.12% | 4 | 0.14% |
| 75 ～ | 男 | 242 | 174 | 71.90% | 39 | 16.12% | 15 | 6.20% | 0 | | 13 | 5.37% | 1 | 0.41% |
| | 女 | 249 | 210 | 84.34% | 19 | 7.63% | 8 | 3.21% | 0 | | 9 | 3.61% | 3 | 1.20% |
| | 計 | 491 | 384 | 78.21% | 58 | 11.81% | 23 | 4.68% | 0 | | 22 | 4.48% | 4 | 0.81% |
| 合計 | | 3452 | 2770 | 80.24% | 348 | 10.08% | 154 | 4.46% | 0 | | 172 | 4.98% | 8 | 0.23% |

5. 脂質検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|--------|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 1 | 33.33% | 1 | 33.33% | 0 | | 1 | 33.33% | 0 | | 0 | |
| | 女 | 31 | 19 | 61.29% | 5 | 16.13% | 6 | 19.35% | 0 | | 1 | 3.23% | 0 | |
| | 計 | 34 | 20 | 58.82% | 6 | 17.65% | 6 | 17.65% | 1 | 2.94% | 1 | 2.94% | 0 | |
| 40 ～ 74 | 男 | 804 | 219 | 27.24% | 160 | 19.90% | 233 | 28.98% | 36 | 4.48% | 13 | 1.62% | 143 | 17.79% |
| | 女 | 2123 | 659 | 31.04% | 438 | 20.63% | 552 | 26.00% | 112 | 5.28% | 34 | 1.60% | 328 | 15.45% |
| | 計 | 2927 | 878 | 30.00% | 598 | 20.43% | 785 | 26.82% | 148 | 5.06% | 47 | 1.61% | 471 | 16.09% |
| 75 ～ | 男 | 242 | 86 | 35.54% | 46 | 19.01% | 46 | 19.01% | 3 | 1.24% | 5 | 2.07% | 56 | 23.14% |
| | 女 | 249 | 75 | 30.12% | 52 | 20.88% | 47 | 18.88% | 6 | 2.41% | 2 | 0.80% | 67 | 26.91% |
| | 計 | 491 | 161 | 32.79% | 98 | 19.96% | 93 | 18.94% | 9 | 1.83% | 7 | 1.43% | 123 | 25.05% |
| 合計 | | 3452 | 1059 | 30.68% | 702 | 20.34% | 884 | 25.61% | 158 | 4.58% | 55 | 1.59% | 594 | 17.21% |

6. 糖代謝検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 1 | 33.33% | 0 | | 1 | 33.33% | 0 | | 0 | | 1 | 33.33% |
| | 女 | 31 | 26 | 83.87% | 4 | 12.90% | 1 | 3.23% | 0 | | 0 | | 0 | |
| | 計 | 34 | 27 | 79.41% | 4 | 11.76% | 2 | 5.88% | 0 | | 0 | | 1 | 2.94% |
| 40 ～ 74 | 男 | 804 | 166 | 20.65% | 247 | 30.72% | 241 | 29.98% | 0 | | 68 | 8.46% | 82 | 10.20% |
| | 女 | 2123 | 782 | 36.83% | 638 | 30.05% | 546 | 25.72% | 0 | | 78 | 3.67% | 79 | 3.72% |
| | 計 | 2927 | 948 | 32.39% | 885 | 30.24% | 787 | 26.89% | 0 | | 146 | 4.99% | 161 | 5.50% |
| 75 ～ | 男 | 242 | 32 | 13.22% | 74 | 30.58% | 81 | 33.47% | 0 | | 18 | 7.44% | 37 | 15.29% |
| | 女 | 249 | 32 | 12.85% | 87 | 34.94% | 82 | 32.93% | 0 | | 18 | 7.23% | 30 | 12.05% |
| | 計 | 491 | 64 | 13.03% | 161 | 32.79% | 163 | 33.20% | 0 | | 36 | 7.33% | 67 | 13.65% |
| 合計 | | 3452 | 1039 | 30.10% | 1050 | 30.42% | 952 | 27.58% | 0 | | 182 | 5.27% | 229 | 6.63% |

7. 腎機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 2 | 66.67% | 0 | | 0 | | 0 | | 1 | 33.33% | 0 | |
| | 女 | 31 | 28 | 90.32% | 3 | 9.68% | 0 | | 0 | | 0 | | 0 | |
| | 計 | 34 | 30 | 88.24% | 3 | 8.82% | 0 | | 0 | | 1 | 2.94% | 0 | |
| 40 ～ 74 | 男 | 804 | 613 | 76.24% | 93 | 11.57% | 72 | 8.96% | 0 | | 26 | 3.23% | 0 | |
| | 女 | 2123 | 1624 | 76.50% | 337 | 15.87% | 150 | 7.07% | 0 | | 12 | 0.57% | 0 | |
| | 計 | 2927 | 2237 | 76.43% | 430 | 14.69% | 222 | 7.58% | 0 | | 38 | 1.30% | 0 | |
| 75 ～ | 男 | 242 | 170 | 70.25% | 32 | 13.22% | 30 | 12.40% | 0 | | 10 | 4.13% | 0 | |
| | 女 | 249 | 131 | 52.61% | 54 | 21.69% | 49 | 19.68% | 0 | | 15 | 6.02% | 0 | |
| | 計 | 491 | 301 | 61.30% | 86 | 17.52% | 79 | 16.09% | 0 | | 25 | 5.09% | 0 | |
| 合計 | | 3452 | 2568 | 74.39% | 519 | 15.03% | 301 | 8.72% | 0 | | 64 | 1.85% | 0 | |

8. 尿酸検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|-------|-------|----|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 3 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 6 | 6 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 9 | 9 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 780 | 623 | 79.87% | 61 | 7.82% | 52 | 6.67% | 4 | 0.51% | 0 | | 40 | 5.13% |
| | 女 | 1362 | 1341 | 98.46% | 12 | 0.88% | 7 | 0.51% | 0 | | 0 | | 2 | 0.15% |
| | 計 | 2142 | 1964 | 91.69% | 73 | 3.41% | 59 | 2.75% | 4 | 0.19% | 0 | | 42 | 1.96% |
| 75 ～ | 男 | 242 | 198 | 81.82% | 20 | 8.26% | 13 | 5.37% | 1 | 0.41% | 0 | | 10 | 4.13% |
| | 女 | 249 | 240 | 96.39% | 6 | 2.41% | 2 | 0.80% | 0 | | 0 | | 1 | 0.40% |
| | 計 | 491 | 438 | 89.21% | 26 | 5.30% | 15 | 3.05% | 1 | 0.20% | 0 | | 11 | 2.24% |
| 合計 | | 2642 | 2411 | 91.26% | 99 | 3.75% | 74 | 2.80% | 5 | 0.19% | 0 | | 53 | 2.01% |

9. 貧血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|--------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 2 | 66.67% | 1 | 33.33% | 0 | | 0 | | 0 | | 0 | |
| | 女 | 9 | 5 | 55.56% | 0 | | 0 | | 4 | 44.44% | 0 | | 0 | |
| | 計 | 12 | 7 | 58.33% | 1 | 8.33% | 0 | | 4 | 33.33% | 0 | | 0 | |
| 40 ～ 74 | 男 | 781 | 644 | 82.46% | 38 | 4.87% | 70 | 8.96% | 18 | 2.30% | 7 | 0.90% | 4 | 0.51% |
| | 女 | 1557 | 1176 | 75.53% | 129 | 8.29% | 172 | 11.05% | 64 | 4.11% | 3 | 0.19% | 13 | 0.83% |
| | 計 | 2338 | 1820 | 77.84% | 167 | 7.14% | 242 | 10.35% | 82 | 3.51% | 10 | 0.43% | 17 | 0.73% |
| 75 ～ | 男 | 242 | 169 | 69.83% | 3 | 1.24% | 49 | 20.25% | 17 | 7.02% | 2 | 0.83% | 2 | 0.83% |
| | 女 | 249 | 168 | 67.47% | 20 | 8.03% | 43 | 17.27% | 15 | 6.02% | 1 | 0.40% | 2 | 0.80% |
| | 計 | 491 | 337 | 68.64% | 23 | 4.68% | 92 | 18.74% | 32 | 6.52% | 3 | 0.61% | 4 | 0.81% |
| 合計 | | 2841 | 2164 | 76.17% | 191 | 6.72% | 334 | 11.76% | 118 | 4.15% | 13 | 0.46% | 21 | 0.74% |

10. 血液一般検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 5 | 5 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 5 | 5 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 6 | 6 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

11. 肝炎検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|----|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 5 | 4 | 80.00% | 0 | | 0 | | 0 | | 1 | 20.00% | 0 | |
| | 計 | 5 | 4 | 80.00% | 0 | | 0 | | 0 | | 1 | 20.00% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5 | 4 | 80.00% | 0 | | 0 | | 0 | | 1 | 20.00% | 0 | |

12. 前立腺がん検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|--------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 7 | 6 | 85.71% | 0 | 0 | 0 | 0 | 0 | 1 | 14.29% | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 7 | 6 | 85.71% | 0 | 0 | 0 | 0 | 0 | 1 | 14.29% | 0 | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 8 | 7 | 87.50% | 0 | 0 | 0 | 0 | 0 | 1 | 12.50% | 0 | 0 | 0 |

13. 便潜血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 8 | 8 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 8 | 8 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 75 ～ | 男 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 9 | 9 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

14. 心電図検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 3 | 3 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 7 | 6 | 85.71% | 0 | 0 | 1 | 14.29% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 10 | 9 | 90.00% | 0 | 0 | 1 | 10.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 787 | 547 | 69.50% | 76 | 9.66% | 96 | 12.20% | 6 | 0.76% | 24 | 3.05% | 38 | 4.83% |
| | 女 | 1469 | 1149 | 78.22% | 115 | 7.83% | 148 | 10.07% | 3 | 0.20% | 31 | 2.11% | 23 | 1.57% |
| | 計 | 2256 | 1696 | 75.18% | 191 | 8.47% | 244 | 10.82% | 9 | 0.40% | 55 | 2.44% | 61 | 2.70% |
| 75 ～ | 男 | 242 | 121 | 50.00% | 27 | 11.16% | 45 | 18.60% | 6 | 2.48% | 20 | 8.26% | 23 | 9.50% |
| | 女 | 249 | 164 | 65.86% | 21 | 8.43% | 30 | 12.05% | 2 | 0.80% | 18 | 7.23% | 14 | 5.62% |
| | 計 | 491 | 285 | 58.04% | 48 | 9.78% | 75 | 15.27% | 8 | 1.63% | 38 | 7.74% | 37 | 7.54% |
| 合計 | | 2757 | 1990 | 72.18% | 239 | 8.67% | 320 | 11.61% | 17 | 0.62% | 93 | 3.37% | 98 | 3.55% |

15. 眼底検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|-------|-------|--------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 247 | 150 | 60.73% | 32 | 12.96% | 28 | 11.34% | 0 | | 23 | 9.31% | 14 | 5.67% |
| | 女 | 378 | 238 | 62.96% | 35 | 9.26% | 48 | 12.70% | 2 | 0.53% | 21 | 5.56% | 34 | 8.99% |
| | 計 | 625 | 388 | 62.08% | 67 | 10.72% | 76 | 12.16% | 2 | 0.32% | 44 | 7.04% | 48 | 7.68% |
| 75 ～ | 男 | 83 | 31 | 37.35% | 8 | 9.64% | 16 | 19.28% | 0 | | 15 | 18.07% | 13 | 15.66% |
| | 女 | 89 | 34 | 38.20% | 8 | 8.99% | 21 | 23.60% | 0 | | 12 | 13.48% | 14 | 15.73% |
| | 計 | 172 | 65 | 37.79% | 16 | 9.30% | 37 | 21.51% | 0 | | 27 | 15.70% | 27 | 15.70% |
| 合計 | | 799 | 455 | 56.95% | 83 | 10.39% | 113 | 14.14% | 2 | 0.25% | 71 | 8.89% | 75 | 9.39% |

16. 眼圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 4 | 4 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 5 | 5 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

17. 肺機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

18. 胸部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 10 | 10 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 12 | 12 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 12 | 12 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

19. 胃部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

20. 胃管内視鏡検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|----|------|----|---------------|---------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |

21. 腹部超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|---------|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 35 | 3 | 8.57% | 1 | 2.86% | 26 | 74.29% | 0 | | 5 | 14.29% | 0 | |
| | 女 | 57 | 8 | 14.04% | 0 | | 46 | 80.70% | 0 | | 3 | 5.26% | 0 | |
| | 計 | 92 | 11 | 11.96% | 1 | 1.09% | 72 | 78.26% | 0 | | 8 | 8.70% | 0 | |
| 75 ～ | 男 | 3 | 0 | | 0 | | 3 | 100.00% | 0 | | 0 | | 0 | |
| | 女 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 5 | 0 | | 0 | | 5 | 100.00% | 0 | | 0 | | 0 | |
| 合計 | | 98 | 11 | 11.22% | 1 | 1.02% | 78 | 79.59% | 0 | | 8 | 8.16% | 0 | |

22. 乳房マンモグラフィ検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|-------|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 81 | 65 | 80.25% | 0 | | 5 | 6.17% | 0 | | 11 | 13.58% | 0 | |
| | 計 | 81 | 65 | 80.25% | 0 | | 5 | 6.17% | 0 | | 11 | 13.58% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 83 | 67 | 80.72% | 0 | | 5 | 6.02% | 0 | | 11 | 13.25% | 0 | |

23. 乳房超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 2 | 1 | 50.00% | 0 | | 1 | 50.00% | 0 | | 0 | | 0 | |
| | 計 | 2 | 1 | 50.00% | 0 | | 1 | 50.00% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 100 | 56 | 56.00% | 0 | | 42 | 42.00% | 0 | | 2 | 2.00% | 0 | |
| | 計 | 100 | 56 | 56.00% | 0 | | 42 | 42.00% | 0 | | 2 | 2.00% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 3 | 2 | 66.67% | 0 | | 1 | 33.33% | 0 | | 0 | | 0 | |
| | 計 | 3 | 2 | 66.67% | 0 | | 1 | 33.33% | 0 | | 0 | | 0 | |
| 合計 | | 105 | 59 | 56.19% | 0 | | 44 | 41.90% | 0 | | 2 | 1.90% | 0 | |

24. 婦人科検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 2 | 1 | 50.00% | 0 | | 0 | | 0 | | 0 | | 1 | 50.00% |
| | 計 | 2 | 1 | 50.00% | 0 | | 0 | | 0 | | 0 | | 1 | 50.00% |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 75 | 68 | 90.67% | 2 | 2.67% | 1 | 1.33% | 0 | | 4 | 5.33% | 0 | |
| | 計 | 75 | 68 | 90.67% | 2 | 2.67% | 1 | 1.33% | 0 | | 4 | 5.33% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 78 | 70 | 89.74% | 2 | 2.56% | 1 | 1.28% | 0 | | 4 | 5.13% | 1 | 1.28% |

④定期健康診断

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|---------|---------------|--------|-----|-------|-------|--------|-----|---------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 1009 | 189 | 18.73% | 198 | 19.62% | 429 | 42.52% | 27 | 2.68% | 152 | 15.06% | 14 | 1.39% |
| | 女 | 799 | 217 | 27.16% | 82 | 10.26% | 386 | 48.31% | 43 | 5.38% | 60 | 7.51% | 11 | 1.38% |
| | 計 | 1808 | 406 | 22.46% | 280 | 15.49% | 815 | 45.08% | 70 | 3.87% | 212 | 11.73% | 25 | 1.38% |
| 40 ～ 74 | 男 | 671 | 38 | 5.66% | 190 | 28.32% | 228 | 33.98% | 23 | 3.43% | 151 | 22.50% | 190 | 28.32% |
| | 女 | 630 | 44 | 6.98% | 101 | 16.03% | 291 | 46.19% | 52 | 8.25% | 81 | 12.86% | 101 | 16.03% |
| | 計 | 1301 | 82 | 6.30% | 291 | 22.37% | 519 | 39.89% | 75 | 5.76% | 232 | 17.83% | 291 | 22.37% |
| 75 ～ | 男 | 5 | 0 | | 3 | 60.00% | 1 | 20.00% | 0 | | 1 | 20.00% | 3 | 60.00% |
| | 女 | 6 | 0 | | 6 | 100.00% | 0 | | 0 | | 0 | | 6 | 100.00% |
| | 計 | 11 | 0 | | 9 | 81.82% | 1 | 9.09% | 0 | | 1 | 9.09% | 9 | 81.82% |
| 合計 | | 3120 | 488 | 15.64% | 580 | 18.59% | 1335 | 42.79% | 145 | 4.65% | 445 | 14.26% | 325 | 10.42% |

1. BMI

| 年代 | 性別 | 受診者数 | ～ 18.4 | | 18.5 ～ 24.9 | | 25.0 ～ | |
|---------|----|------|--------|--------|-------------|--------|--------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 1003 | 82 | 8.18% | 630 | 62.81% | 291 | 29.01% |
| | 女 | 799 | 151 | 18.90% | 519 | 64.96% | 129 | 16.15% |
| | 計 | 1802 | 233 | 12.93% | 1149 | 63.76% | 420 | 23.31% |
| 40 ～ 74 | 男 | 658 | 30 | 4.56% | 361 | 54.86% | 267 | 40.58% |
| | 女 | 629 | 69 | 10.97% | 423 | 67.25% | 137 | 21.78% |
| | 計 | 1287 | 99 | 7.69% | 784 | 60.92% | 404 | 31.39% |
| 75 ～ | 男 | 5 | 0 | | 2 | 40.00% | 3 | 60.00% |
| | 女 | 6 | 1 | 16.67% | 4 | 66.67% | 1 | 16.67% |
| | 計 | 11 | 1 | 9.09% | 6 | 54.55% | 4 | 36.36% |
| 合計 | | 3100 | 333 | 10.74% | 1939 | 62.55% | 828 | 26.71% |

2. 血圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|--------|-----|---------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 1009 | 721 | 71.46% | 181 | 17.94% | 85 | 8.42% | 0 | | 15 | 1.49% | 7 | 0.69% |
| | 女 | 799 | 720 | 90.11% | 36 | 4.51% | 28 | 3.50% | 0 | | 10 | 1.25% | 5 | 0.63% |
| | 計 | 1808 | 1441 | 79.70% | 217 | 12.00% | 113 | 6.25% | 0 | | 25 | 1.38% | 12 | 0.66% |
| 40 ～ 74 | 男 | 669 | 232 | 34.68% | 100 | 14.95% | 118 | 17.64% | 0 | | 65 | 9.72% | 154 | 23.02% |
| | 女 | 629 | 377 | 59.94% | 74 | 11.76% | 83 | 13.20% | 0 | | 26 | 4.13% | 69 | 10.97% |
| | 計 | 1298 | 609 | 46.92% | 174 | 13.41% | 201 | 15.49% | 0 | | 91 | 7.01% | 223 | 17.18% |
| 75 ～ | 男 | 5 | 1 | 20.00% | 0 | | 0 | | 0 | | 1 | 20.00% | 3 | 60.00% |
| | 女 | 6 | 0 | | 0 | | 0 | | 0 | | 0 | | 6 | 100.00% |
| | 計 | 11 | 1 | 9.09% | 0 | | 0 | | 0 | | 1 | 9.09% | 9 | 81.82% |
| 合計 | | 3117 | 2051 | 65.80% | 391 | 12.54% | 314 | 10.07% | 0 | | 117 | 3.75% | 244 | 7.83% |

3. 聴力検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 789 | 782 | 99.11% | 0 | | 7 | 0.89% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 789 | 782 | 99.11% | 0 | | 7 | 0.89% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 622 | 583 | 93.73% | 0 | | 39 | 6.27% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 622 | 583 | 93.73% | 0 | | 39 | 6.27% | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 4 | 2 | 50.00% | 0 | | 2 | 50.00% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 4 | 2 | 50.00% | 0 | | 2 | 50.00% | 0 | | 0 | | 0 | |
| 合計 | | 1415 | 1367 | 96.61% | 0 | | 48 | 3.39% | 0 | | 0 | | 0 | |

4. 肝機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 577 | 543 | 94.11% | 16 | 2.77% | 6 | 1.04% | 0 | | 12 | 2.08% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 577 | 543 | 94.11% | 16 | 2.77% | 6 | 1.04% | 0 | | 12 | 2.08% | 0 | |
| 40 ～ 74 | 男 | 566 | 507 | 89.58% | 26 | 4.59% | 17 | 3.00% | 0 | | 16 | 2.83% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 566 | 507 | 89.58% | 26 | 4.59% | 17 | 3.00% | 0 | | 16 | 2.83% | 0 | |
| 75 ～ | 男 | 6 | 4 | 66.67% | 1 | 16.67% | 1 | 16.67% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 4 | 66.67% | 1 | 16.67% | 1 | 16.67% | 0 | | 0 | | 0 | |
| 合計 | | 1149 | 1054 | 91.73% | 43 | 3.74% | 24 | 2.09% | 0 | | 28 | 2.44% | 0 | |

5. 脂質検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 577 | 415 | 71.92% | 78 | 13.52% | 50 | 8.67% | 7 | 1.21% | 24 | 4.16% | 3 | 0.52% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 577 | 415 | 71.92% | 78 | 13.52% | 50 | 8.67% | 7 | 1.21% | 24 | 4.16% | 3 | 0.52% |
| 40 ～ 74 | 男 | 577 | 222 | 38.47% | 128 | 22.18% | 142 | 24.61% | 30 | 5.20% | 10 | 1.73% | 45 | 7.80% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 577 | 222 | 38.47% | 128 | 22.18% | 142 | 24.61% | 30 | 5.20% | 10 | 1.73% | 45 | 7.80% |
| 75 ～ | 男 | 6 | 2 | 33.33% | 0 | | 1 | 16.67% | 0 | | 0 | | 3 | 50.00% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 2 | 33.33% | 0 | | 1 | 16.67% | 0 | | 0 | | 3 | 50.00% |
| 合計 | | 1160 | 639 | 55.09% | 206 | 17.76% | 193 | 16.64% | 37 | 3.19% | 34 | 2.93% | 51 | 4.40% |

6. 糖代謝検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|----|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 793 | 627 | 79.07% | 149 | 18.79% | 13 | 1.64% | 0 | | 2 | 0.25% | 2 | 0.25% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 793 | 627 | 79.07% | 149 | 18.79% | 13 | 1.64% | 0 | | 2 | 0.25% | 2 | 0.25% |
| 40 ～ 74 | 男 | 628 | 283 | 45.06% | 236 | 37.58% | 87 | 13.85% | 0 | | 12 | 1.91% | 10 | 1.59% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 628 | 283 | 45.06% | 236 | 37.58% | 87 | 13.85% | 0 | | 12 | 1.91% | 10 | 1.59% |
| 75 ～ | 男 | 6 | 1 | 16.67% | 0 | | 3 | 50.00% | 0 | | 0 | | 2 | 33.33% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 1 | 16.67% | 0 | | 3 | 50.00% | 0 | | 0 | | 2 | 33.33% |
| 合計 | | 1427 | 911 | 63.84% | 385 | 26.98% | 103 | 7.22% | 0 | | 14 | 0.98% | 14 | 0.98% |

7. 腎機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 782 | 747 | 95.52% | 22 | 2.81% | 12 | 1.53% | 0 | | 1 | 0.13% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 782 | 747 | 95.52% | 22 | 2.81% | 12 | 1.53% | 0 | | 1 | 0.13% | 0 | |
| 40 ～ 74 | 男 | 626 | 571 | 91.21% | 30 | 4.79% | 22 | 3.51% | 0 | | 3 | 0.48% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 626 | 571 | 91.21% | 30 | 4.79% | 22 | 3.51% | 0 | | 3 | 0.48% | 0 | |
| 75 ～ | 男 | 5 | 3 | 60.00% | 2 | 40.00% | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 5 | 3 | 60.00% | 2 | 40.00% | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1413 | 1321 | 93.49% | 54 | 3.82% | 34 | 2.41% | 0 | | 4 | 0.28% | 0 | |

8. 尿酸検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 72 | 70 | 97.22% | 1 | 1.39% | 1 | 1.39% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 72 | 70 | 97.22% | 1 | 1.39% | 1 | 1.39% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 132 | 131 | 99.24% | 0 | | 1 | 0.76% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 132 | 131 | 99.24% | 0 | | 1 | 0.76% | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 205 | 202 | 98.54% | 1 | 0.49% | 2 | 0.98% | 0 | | 0 | | 0 | |

9. 貧血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 575 | 382 | 66.43% | 64 | 11.13% | 87 | 15.13% | 38 | 6.61% | 3 | 0.52% | 1 | 0.17% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 575 | 382 | 66.43% | 64 | 11.13% | 87 | 15.13% | 38 | 6.61% | 3 | 0.52% | 1 | 0.17% |
| 40 ～ 74 | 男 | 565 | 408 | 72.21% | 47 | 8.32% | 67 | 11.86% | 36 | 6.37% | 2 | 0.35% | 5 | 0.88% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 565 | 408 | 72.21% | 47 | 8.32% | 67 | 11.86% | 36 | 6.37% | 2 | 0.35% | 5 | 0.88% |
| 75 ～ | 男 | 6 | 2 | 33.33% | 0 | | 4 | 66.67% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 2 | 33.33% | 0 | | 4 | 66.67% | 0 | | 0 | | 0 | |
| 合計 | | 1146 | 792 | 69.11% | 111 | 9.69% | 158 | 13.79% | 74 | 6.46% | 5 | 0.44% | 6 | 0.52% |

10. 血液一般検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 62 | 53 | 85.48% | 2 | 3.23% | 3 | 4.84% | 0 | | 4 | 6.45% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 62 | 53 | 85.48% | 2 | 3.23% | 3 | 4.84% | 0 | | 4 | 6.45% | 0 | |
| 40 ～ 74 | 男 | 125 | 112 | 89.60% | 1 | 0.80% | 8 | 6.40% | 0 | | 4 | 3.20% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 125 | 112 | 89.60% | 1 | 0.80% | 8 | 6.40% | 0 | | 4 | 3.20% | 0 | |
| 75 ～ | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 188 | 166 | 88.30% | 3 | 1.60% | 11 | 5.85% | 0 | | 8 | 4.26% | 0 | |

11. 肝炎検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 33 | 33 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 33 | 33 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 21 | 21 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 21 | 21 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 54 | 54 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

12. 前立腺がん検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--|------|--|---------------|--|-----|--|-------|--|-----|--|
| | | | | | | | | | | | | | | |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |

13. 便潜血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--|---------------|--|-----|--|-------|-------|-----|--|
| | | | | | | | | | | | | | | |
| ～ 39 | 男 | 32 | 30 | 93.75% | 0 | | 0 | | 0 | | 2 | 6.25% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 32 | 30 | 93.75% | 0 | | 0 | | 0 | | 2 | 6.25% | 0 | |
| 40 ～ 74 | 男 | 99 | 95 | 95.96% | 0 | | 0 | | 0 | | 4 | 4.04% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 99 | 95 | 95.96% | 0 | | 0 | | 0 | | 4 | 4.04% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 131 | 125 | 95.42% | 0 | | 0 | | 0 | | 6 | 4.58% | 0 | |

14. 心電図検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|--|-------|--------|-----|-------|
| | | | | | | | | | | | | | | |
| ～ 39 | 男 | 534 | 478 | 89.51% | 24 | 4.49% | 28 | 5.24% | 0 | | 4 | 0.75% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 534 | 478 | 89.51% | 24 | 4.49% | 28 | 5.24% | 0 | | 4 | 0.75% | 0 | |
| 40 ～ 74 | 男 | 562 | 483 | 85.94% | 32 | 5.69% | 39 | 6.94% | 0 | | 6 | 1.07% | 2 | 0.36% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 562 | 483 | 85.94% | 32 | 5.69% | 39 | 6.94% | 0 | | 6 | 1.07% | 2 | 0.36% |
| 75 ～ | 男 | 6 | 4 | 66.67% | 0 | | 1 | 16.67% | 0 | | 1 | 16.67% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 4 | 66.67% | 0 | | 1 | 16.67% | 0 | | 1 | 16.67% | 0 | |
| 合計 | | 1102 | 965 | 87.57% | 56 | 5.08% | 68 | 6.17% | 0 | | 11 | 1.00% | 2 | 0.18% |

15. 眼底検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|----|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 4 | 3 | 75.00% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 25.00% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 4 | 3 | 75.00% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 25.00% | 0 | 0 |
| 40 ～ 74 | 男 | 8 | 6 | 75.00% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 25.00% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 8 | 6 | 75.00% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 25.00% | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 合計 | | 12 | 9 | 75.00% | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 25.00% | 0 | 0 |

16. 眼圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|----|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 合計 | | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |

17. 肺機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|----|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |
| 合計 | | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | 0 | 0 |

18. 胸部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|---------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 727 | 659 | 90.65% | 1 | 0.14% | 67 | 9.22% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 727 | 659 | 90.65% | 1 | 0.14% | 67 | 9.22% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 564 | 455 | 80.67% | 0 | | 102 | 18.09% | 0 | 0 | 4 | 0.71% | 3 | 0.53% |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 564 | 455 | 80.67% | 0 | | 102 | 18.09% | 0 | 0 | 4 | 0.71% | 3 | 0.53% |
| 75 ～ | 男 | 5 | 0 | | 0 | | 5 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 5 | 0 | | 0 | | 5 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 1296 | 1114 | 85.96% | 1 | 0.08% | 174 | 13.43% | 0 | 0 | 4 | 0.31% | 3 | 0.23% |

19. 胃部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 7 | 5 | 71.43% | 0 | | 2 | 28.57% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 7 | 5 | 71.43% | 0 | | 2 | 28.57% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 66 | 60 | 90.91% | 0 | | 5 | 7.58% | 0 | 0 | 1 | 1.52% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 66 | 60 | 90.91% | 0 | | 5 | 7.58% | 0 | 0 | 1 | 1.52% | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 73 | 65 | 89.04% | 0 | | 7 | 9.59% | 0 | 0 | 1 | 1.37% | 0 | 0 |

20. 胃部内視鏡検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 6 | 1 | 16.67% | 0 | | 5 | 83.33% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 6 | 1 | 16.67% | 0 | | 5 | 83.33% | 0 | 0 | 0 | 0 | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 6 | 1 | 16.67% | 0 | | 5 | 83.33% | 0 | 0 | 0 | 0 | 0 | 0 |

21. 腹部超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|---------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 19 | 4 | 21.05% | 0 | | 14 | 73.68% | 0 | | 1 | 5.26% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 19 | 4 | 21.05% | 0 | | 14 | 73.68% | 0 | | 1 | 5.26% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 21 | 4 | 19.05% | 0 | | 16 | 76.19% | 0 | | 1 | 4.76% | 0 | |

22. 乳房マンモグラフィ検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 23 | 21 | 91.30% | 0 | | 2 | 8.70% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 23 | 21 | 91.30% | 0 | | 2 | 8.70% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 77 | 61 | 79.22% | 0 | | 11 | 14.29% | 0 | | 5 | 6.49% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 77 | 61 | 79.22% | 0 | | 11 | 14.29% | 0 | | 5 | 6.49% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 100 | 82 | 82.00% | 0 | | 13 | 13.00% | 0 | | 5 | 5.00% | 0 | |

23. 乳房超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 65 | 40 | 61.54% | 0 | | 23 | 35.38% | 0 | | 2 | 3.08% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 65 | 40 | 61.54% | 0 | | 23 | 35.38% | 0 | | 2 | 3.08% | 0 | |
| 40 ～ 74 | 男 | 56 | 25 | 44.64% | 0 | | 30 | 53.57% | 0 | | 1 | 1.79% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 56 | 25 | 44.64% | 0 | | 30 | 53.57% | 0 | | 1 | 1.79% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 121 | 65 | 53.72% | 0 | | 53 | 43.80% | 0 | | 3 | 2.48% | 0 | |

24. 婦人科検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 64 | 64 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 64 | 64 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 111 | 104 | 93.69% | 4 | 3.60% | 1 | 0.90% | 0 | | 2 | 1.80% | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 111 | 104 | 93.69% | 4 | 3.60% | 1 | 0.90% | 0 | | 2 | 1.80% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 175 | 168 | 96.00% | 4 | 2.29% | 1 | 0.57% | 0 | | 2 | 1.14% | 0 | |

⑤その他

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|-------|-------|--------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 105 | 41 | 39.05% | 10 | 9.52% | 32 | 30.48% | 7 | 6.67% | 15 | 14.29% | 0 | |
| | 女 | 586 | 351 | 59.90% | 24 | 4.10% | 171 | 29.18% | 8 | 1.37% | 27 | 4.61% | 5 | 0.85% |
| | 計 | 691 | 392 | 56.73% | 34 | 4.92% | 203 | 29.38% | 15 | 2.17% | 42 | 6.08% | 5 | 0.72% |
| 40 ～ 74 | 男 | 258 | 124 | 48.06% | 6 | 2.33% | 68 | 26.36% | 13 | 5.04% | 41 | 15.89% | 6 | 2.33% |
| | 女 | 1185 | 830 | 70.04% | 17 | 1.43% | 217 | 18.31% | 11 | 0.93% | 97 | 8.19% | 17 | 1.43% |
| | 計 | 1443 | 954 | 66.11% | 23 | 1.59% | 285 | 19.75% | 24 | 1.66% | 138 | 9.56% | 23 | 1.59% |
| 75 ～ | 男 | 10 | 2 | 20.00% | 3 | 30.00% | 2 | 20.00% | 0 | | 3 | 30.00% | 3 | 30.00% |
| | 女 | 39 | 33 | 84.62% | 0 | | 3 | 7.69% | 1 | 2.56% | 2 | 5.13% | 0 | |
| | 計 | 49 | 35 | 71.43% | 3 | 6.12% | 5 | 10.20% | 1 | 2.04% | 5 | 10.20% | 3 | 6.12% |
| 合計 | | 2183 | 1381 | 63.26% | 60 | 2.75% | 493 | 22.58% | 40 | 1.83% | 185 | 8.47% | 31 | 1.42% |

1. BMI

| 年代 | 性別 | 受診者数 | ～ 18.4 | | 18.5 ～ 24.9 | | 25.0 ～ | |
|---------|----|------|--------|--------|-------------|---------|--------|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 46 | 3 | 6.52% | 28 | 60.87% | 15 | 32.61% |
| | 女 | 135 | 25 | 18.52% | 87 | 64.44% | 23 | 17.04% |
| | 計 | 181 | 28 | 15.47% | 115 | 63.54% | 38 | 20.99% |
| 40 ～ 74 | 男 | 47 | 1 | 2.13% | 31 | 65.96% | 15 | 31.91% |
| | 女 | 59 | 9 | 15.25% | 37 | 62.71% | 13 | 22.03% |
| | 計 | 106 | 10 | 9.43% | 68 | 64.15% | 28 | 26.42% |
| 75 ～ | 男 | 5 | 0 | | 5 | 100.00% | 0 | |
| | 女 | 1 | 0 | | 1 | 100.00% | 0 | |
| | 計 | 6 | 0 | | 6 | 100.00% | 0 | |
| 合計 | | 293 | 38 | 12.97% | 189 | 64.51% | 66 | 22.53% |

2. 血圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|---------|-----|-------|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 46 | 33 | 71.74% | 11 | 23.91% | 2 | 4.35% | 0 | | 0 | | 0 | |
| | 女 | 136 | 126 | 92.65% | 5 | 3.68% | 4 | 2.94% | 0 | | 0 | | 1 | 0.74% |
| | 計 | 182 | 159 | 87.36% | 16 | 8.79% | 6 | 3.30% | 0 | | 0 | | 1 | 0.55% |
| 40 ～ 74 | 男 | 50 | 22 | 44.00% | 9 | 18.00% | 10 | 20.00% | 2 | 4.00% | 3 | 6.00% | 4 | 8.00% |
| | 女 | 60 | 31 | 51.67% | 8 | 13.33% | 9 | 15.00% | 1 | 1.67% | 2 | 3.33% | 9 | 15.00% |
| | 計 | 110 | 53 | 48.18% | 17 | 15.45% | 19 | 17.27% | 3 | 2.73% | 5 | 4.55% | 13 | 11.82% |
| 75 ～ | 男 | 5 | 0 | | 1 | 20.00% | 2 | 40.00% | 0 | | 0 | | 2 | 40.00% |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 6 | 0 | | 1 | 16.67% | 3 | 50.00% | 0 | | 0 | | 2 | 33.33% |
| 合計 | | 298 | 212 | 71.14% | 34 | 11.41% | 28 | 9.40% | 3 | 1.01% | 5 | 1.68% | 16 | 5.37% |

3. 聴力検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|--------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 46 | 44 | 95.65% | 0 | 0 | 2 | 4.35% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 131 | 129 | 98.47% | 0 | 0 | 2 | 1.53% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 177 | 173 | 97.74% | 0 | 0 | 4 | 2.26% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 40 | 32 | 80.00% | 0 | 0 | 8 | 20.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 35 | 34 | 97.14% | 0 | 0 | 1 | 2.86% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 75 | 66 | 88.00% | 0 | 0 | 9 | 12.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| 75 ～ | 男 | 5 | 1 | 20.00% | 0 | 0 | 4 | 80.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 6 | 2 | 33.33% | 0 | 0 | 4 | 66.67% | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 258 | 241 | 93.41% | 0 | 0 | 17 | 6.59% | 0 | 0 | 0 | 0 | 0 | 0 |

4. 肝機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|-------|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 57 | 38 | 66.67% | 4 | 7.02% | 6 | 10.53% | 2 | 3.51% | 7 | 12.28% | 0 | 0 |
| | 女 | 117 | 111 | 94.87% | 3 | 2.56% | 2 | 1.71% | 0 | 0 | 1 | 0.85% | 0 | 0 |
| | 計 | 174 | 149 | 85.63% | 7 | 4.02% | 8 | 4.60% | 2 | 1.15% | 8 | 4.60% | 0 | 0 |
| 40 ～ 74 | 男 | 61 | 31 | 50.82% | 15 | 24.59% | 7 | 11.48% | 2 | 3.28% | 6 | 9.84% | 0 | 0 |
| | 女 | 69 | 55 | 79.71% | 8 | 11.59% | 2 | 2.90% | 0 | 0 | 4 | 5.80% | 0 | 0 |
| | 計 | 130 | 86 | 66.15% | 23 | 17.69% | 9 | 6.92% | 2 | 1.54% | 10 | 7.69% | 0 | 0 |
| 75 ～ | 男 | 5 | 3 | 60.00% | 2 | 40.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 6 | 4 | 66.67% | 2 | 33.33% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 310 | 239 | 77.10% | 32 | 10.32% | 17 | 5.48% | 4 | 1.29% | 18 | 5.81% | 0 | 0 |

5. 脂質検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|--------|-----|---------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 57 | 22 | 38.60% | 8 | 14.04% | 18 | 31.58% | 6 | 10.53% | 3 | 5.26% | 0 | 0 |
| | 女 | 117 | 80 | 68.38% | 18 | 15.38% | 10 | 8.55% | 4 | 3.42% | 5 | 4.27% | 0 | 0 |
| | 計 | 174 | 102 | 58.62% | 26 | 14.94% | 28 | 16.09% | 10 | 5.75% | 8 | 4.60% | 0 | 0 |
| 40 ～ 74 | 男 | 54 | 10 | 18.52% | 13 | 24.07% | 20 | 37.04% | 6 | 11.11% | 0 | 0 | 5 | 9.26% |
| | 女 | 73 | 31 | 42.47% | 10 | 13.70% | 20 | 27.40% | 4 | 5.48% | 1 | 1.37% | 7 | 9.59% |
| | 計 | 127 | 41 | 32.28% | 23 | 18.11% | 40 | 31.50% | 10 | 7.87% | 1 | 0.79% | 12 | 9.45% |
| 75 ～ | 男 | 5 | 1 | 20.00% | 1 | 20.00% | 2 | 40.00% | 1 | 20.00% | 0 | 0 | 0 | 0 |
| | 女 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 100.00% | 0 | 0 | 0 | 0 |
| | 計 | 6 | 1 | 16.67% | 1 | 16.67% | 2 | 33.33% | 2 | 33.33% | 0 | 0 | 0 | 0 |
| 合計 | | 307 | 144 | 46.91% | 50 | 16.29% | 70 | 22.80% | 22 | 7.17% | 9 | 2.93% | 12 | 3.91% |

6. 糖代謝検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|---------|---------------|--------|-----|-------|-------|-------|-----|--------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 59 | 48 | 81.36% | 8 | 13.56% | 2 | 3.39% | 0 | | 1 | 1.69% | 0 | |
| | 女 | 141 | 113 | 80.14% | 19 | 13.48% | 4 | 2.84% | 2 | 1.42% | 2 | 1.42% | 1 | 0.71% |
| | 計 | 200 | 161 | 80.50% | 27 | 13.50% | 6 | 3.00% | 2 | 1.00% | 3 | 1.50% | 1 | 0.50% |
| 40 ～ 74 | 男 | 58 | 23 | 39.66% | 20 | 34.48% | 11 | 18.97% | 1 | 1.72% | 1 | 1.72% | 2 | 3.45% |
| | 女 | 76 | 45 | 59.21% | 20 | 26.32% | 7 | 9.21% | 1 | 1.32% | 3 | 3.95% | 0 | |
| | 計 | 134 | 68 | 50.75% | 40 | 29.85% | 18 | 13.43% | 2 | 1.49% | 4 | 2.99% | 2 | 1.49% |
| 75 ～ | 男 | 5 | 3 | 60.00% | 1 | 20.00% | 0 | | 0 | | 0 | | 1 | 20.00% |
| | 女 | 1 | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 3 | 50.00% | 2 | 33.33% | 0 | | 0 | | 0 | | 1 | 16.67% |
| 合計 | | 340 | 232 | 68.24% | 69 | 20.29% | 24 | 7.06% | 4 | 1.18% | 7 | 2.06% | 4 | 1.18% |

7. 腎機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|-------|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 51 | 43 | 84.31% | 8 | 15.69% | 0 | | 0 | | 0 | | 0 | |
| | 女 | 142 | 121 | 85.21% | 9 | 6.34% | 11 | 7.75% | 0 | | 1 | 0.70% | 0 | |
| | 計 | 193 | 164 | 84.97% | 17 | 8.81% | 11 | 5.70% | 0 | | 1 | 0.52% | 0 | |
| 40 ～ 74 | 男 | 55 | 43 | 78.18% | 4 | 7.27% | 7 | 12.73% | 1 | 1.82% | 0 | | 0 | |
| | 女 | 71 | 50 | 70.42% | 13 | 18.31% | 7 | 9.86% | 0 | | 1 | 1.41% | 0 | |
| | 計 | 126 | 93 | 73.81% | 17 | 13.49% | 14 | 11.11% | 1 | 0.79% | 1 | 0.79% | 0 | |
| 75 ～ | 男 | 5 | 3 | 60.00% | 1 | 20.00% | 1 | 20.00% | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 4 | 66.67% | 1 | 16.67% | 1 | 16.67% | 0 | | 0 | | 0 | |
| 合計 | | 325 | 261 | 80.31% | 35 | 10.77% | 26 | 8.00% | 1 | 0.31% | 2 | 0.62% | 0 | |

8. 尿酸検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|-------|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 40 | 33 | 82.50% | 3 | 7.50% | 3 | 7.50% | 1 | 2.50% | 0 | | 0 | |
| | 女 | 45 | 43 | 95.56% | 1 | 2.22% | 1 | 2.22% | 0 | | 0 | | 0 | |
| | 計 | 85 | 76 | 89.41% | 4 | 4.71% | 4 | 4.71% | 1 | 1.18% | 0 | | 0 | |
| 40 ～ 74 | 男 | 46 | 36 | 78.26% | 7 | 15.22% | 2 | 4.35% | 1 | 2.17% | 0 | | 0 | |
| | 女 | 50 | 49 | 98.00% | 1 | 2.00% | 0 | | 0 | | 0 | | 0 | |
| | 計 | 96 | 85 | 88.54% | 8 | 8.33% | 2 | 2.08% | 1 | 1.04% | 0 | | 0 | |
| 75 ～ | 男 | 5 | 4 | 80.00% | 0 | | 1 | 20.00% | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 5 | 83.33% | 0 | | 1 | 16.67% | 0 | | 0 | | 0 | |
| 合計 | | 187 | 166 | 88.77% | 12 | 6.42% | 7 | 3.74% | 2 | 1.07% | 0 | | 0 | |

9. 貧血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|--------|---------------|---------|-----|-------|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 57 | 38 | 66.67% | 14 | 24.56% | 1 | 1.75% | 2 | 3.51% | 2 | 3.51% | 0 | |
| | 女 | 120 | 80 | 66.67% | 21 | 17.50% | 12 | 10.00% | 5 | 4.17% | 1 | 0.83% | 1 | 0.83% |
| | 計 | 177 | 118 | 66.67% | 35 | 19.77% | 13 | 7.34% | 7 | 3.95% | 3 | 1.69% | 1 | 0.56% |
| 40 ～ 74 | 男 | 61 | 49 | 80.33% | 7 | 11.48% | 3 | 4.92% | 2 | 3.28% | 0 | | 0 | |
| | 女 | 72 | 54 | 75.00% | 4 | 5.56% | 8 | 11.11% | 3 | 4.17% | 1 | 1.39% | 2 | 2.78% |
| | 計 | 133 | 103 | 77.44% | 11 | 8.27% | 11 | 8.27% | 5 | 3.76% | 1 | 0.75% | 2 | 1.50% |
| 75 ～ | 男 | 5 | 3 | 60.00% | 0 | | 2 | 40.00% | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 6 | 3 | 50.00% | 0 | | 3 | 50.00% | 0 | | 0 | | 0 | |
| 合計 | | 316 | 224 | 70.89% | 46 | 14.56% | 27 | 8.54% | 12 | 3.80% | 4 | 1.27% | 3 | 0.95% |

10. 血液一般検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|--------|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 41 | 39 | 95.12% | 0 | | 0 | | 0 | | 2 | 4.88% | 0 | |
| | 女 | 43 | 36 | 83.72% | 3 | 6.98% | 0 | | 0 | | 4 | 9.30% | 0 | |
| | 計 | 84 | 75 | 89.29% | 3 | 3.57% | 0 | | 0 | | 6 | 7.14% | 0 | |
| 40 ～ 74 | 男 | 38 | 30 | 78.95% | 2 | 5.26% | 0 | | 2 | 5.26% | 4 | 10.53% | 0 | |
| | 女 | 24 | 19 | 79.17% | 0 | | 1 | 4.17% | 3 | 12.50% | 1 | 4.17% | 0 | |
| | 計 | 62 | 49 | 79.03% | 2 | 3.23% | 1 | 1.61% | 5 | 8.06% | 5 | 8.06% | 0 | |
| 75 ～ | 男 | 5 | 5 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 6 | 6 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 152 | 130 | 85.53% | 5 | 3.29% | 1 | 0.66% | 5 | 3.29% | 11 | 7.24% | 0 | |

11. 肝炎検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 17 | 17 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 7 | 7 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 24 | 24 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 14 | 14 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 15 | 15 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 29 | 29 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 53 | 53 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

12. 前立腺がん検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 4 | 4 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 4 | 4 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 57 | 55 | 96.49% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 3.51% | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 57 | 55 | 96.49% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 3.51% | 0 | 0 |
| 75 ～ | 男 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 61 | 59 | 96.72% | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 3.28% | 0 | 0 |

13. 便潜血検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 34 | 33 | 97.06% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.94% | 0 | 0 |
| | 女 | 51 | 51 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 85 | 84 | 98.82% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1.18% | 0 | 0 |
| 40 ～ 74 | 男 | 47 | 41 | 87.23% | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 12.77% | 0 | 0 |
| | 女 | 64 | 64 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 111 | 105 | 94.59% | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 5.41% | 0 | 0 |
| 75 ～ | 男 | 5 | 4 | 80.00% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 20.00% | 0 | 0 |
| | 女 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 6 | 5 | 83.33% | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 16.67% | 0 | 0 |
| 合計 | | 202 | 194 | 96.04% | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 3.96% | 0 | 0 |

14. 心電図検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|--------|---------------|--------|-----|----|-------|-------|-----|-------|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 45 | 43 | 95.56% | 1 | 2.22% | 1 | 2.22% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 101 | 84 | 83.17% | 11 | 10.89% | 6 | 5.94% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 146 | 127 | 86.99% | 12 | 8.22% | 7 | 4.79% | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 ～ 74 | 男 | 46 | 39 | 84.78% | 3 | 6.52% | 2 | 4.35% | 0 | 0 | 1 | 2.17% | 1 | 2.17% |
| | 女 | 62 | 42 | 67.74% | 12 | 19.35% | 4 | 6.45% | 0 | 0 | 3 | 4.84% | 1 | 1.61% |
| | 計 | 108 | 81 | 75.00% | 15 | 13.89% | 6 | 5.56% | 0 | 0 | 4 | 3.70% | 2 | 1.85% |
| 75 ～ | 男 | 5 | 3 | 60.00% | 1 | 20.00% | 1 | 20.00% | 0 | 0 | 0 | 0 | 0 | 0 |
| | 女 | 1 | 1 | 100.00% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 計 | 6 | 4 | 66.67% | 1 | 16.67% | 1 | 16.67% | 0 | 0 | 0 | 0 | 0 | 0 |
| 合計 | | 260 | 212 | 81.54% | 28 | 10.77% | 14 | 5.38% | 0 | 0 | 4 | 1.54% | 2 | 0.77% |

15. 眼底検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|---------|-----|----|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 5 | 4 | 80.00% | 0 | | 0 | | 0 | | 1 | 20.00% | 0 | |
| | 女 | 4 | 3 | 75.00% | 0 | | 1 | 25.00% | 0 | | 0 | | 0 | |
| | 計 | 9 | 7 | 77.78% | 0 | | 1 | 11.11% | 0 | | 1 | 11.11% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| 合計 | | 11 | 8 | 72.73% | 0 | | 2 | 18.18% | 0 | | 1 | 9.09% | 0 | |

16. 眼圧検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 2 | 2 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

17. 肺機能検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|----|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |

18. 胸部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|---------|-----|----|-------|----|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 44 | 41 | 93.18% | 0 | | 3 | 6.82% | 0 | | 0 | | 0 | |
| | 女 | 133 | 122 | 91.73% | 0 | | 11 | 8.27% | 0 | | 0 | | 0 | |
| | 計 | 177 | 163 | 92.09% | 0 | | 14 | 7.91% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 46 | 40 | 86.96% | 0 | | 6 | 13.04% | 0 | | 0 | | 0 | |
| | 女 | 74 | 61 | 82.43% | 0 | | 13 | 17.57% | 0 | | 0 | | 0 | |
| | 計 | 120 | 101 | 84.17% | 0 | | 19 | 15.83% | 0 | | 0 | | 0 | |
| 75 ～ | 男 | 5 | 0 | | 0 | | 5 | 100.00% | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 6 | 0 | | 0 | | 6 | 100.00% | 0 | | 0 | | 0 | |
| 合計 | | 303 | 264 | 87.13% | 0 | | 39 | 12.87% | 0 | | 0 | | 0 | |

19. 胃部X線検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|---------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 34 | 29 | 85.29% | 0 | | 3 | 8.82% | 0 | | 2 | 5.88% | 0 | |
| | 女 | 24 | 20 | 83.33% | 0 | | 3 | 12.50% | 0 | | 1 | 4.17% | 0 | |
| | 計 | 58 | 49 | 84.48% | 0 | | 6 | 10.34% | 0 | | 3 | 5.17% | 0 | |
| 40 ～ 74 | 男 | 32 | 26 | 81.25% | 0 | | 3 | 9.38% | 0 | | 3 | 9.38% | 0 | |
| | 女 | 24 | 22 | 91.67% | 0 | | 2 | 8.33% | 0 | | 0 | | 0 | |
| | 計 | 56 | 48 | 85.71% | 0 | | 5 | 8.93% | 0 | | 3 | 5.36% | 0 | |
| 75 ～ | 男 | 3 | 2 | 66.67% | 0 | | 1 | 33.33% | 0 | | 0 | | 0 | |
| | 女 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 4 | 2 | 50.00% | 0 | | 2 | 50.00% | 0 | | 0 | | 0 | |
| 合計 | | 118 | 99 | 83.90% | 0 | | 13 | 11.02% | 0 | | 6 | 5.08% | 0 | |

20. 胃部内視鏡検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|---------|-----|-------|-------|--------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 1 | 0 | | 0 | | 1 | 100.00% | 0 | | 0 | | 0 | |
| | 女 | 3 | 1 | 33.33% | 0 | | 2 | 66.67% | 0 | | 0 | | 0 | |
| | 計 | 4 | 1 | 25.00% | 0 | | 3 | 75.00% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 11 | 1 | 9.09% | 0 | | 6 | 54.55% | 1 | 9.09% | 3 | 27.27% | 0 | |
| | 女 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| | 計 | 13 | 1 | 7.69% | 0 | | 8 | 61.54% | 1 | 7.69% | 3 | 23.08% | 0 | |
| 75 ～ | 男 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 2 | 0 | | 0 | | 2 | 100.00% | 0 | | 0 | | 0 | |
| 合計 | | 19 | 2 | 10.53% | 0 | | 13 | 68.42% | 1 | 5.26% | 3 | 15.79% | 0 | |

21. 腹部超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|-------|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 5 | 2 | 40.00% | 0 | | 3 | 60.00% | 0 | | 0 | | 0 | |
| | 女 | 12 | 4 | 33.33% | 1 | 8.33% | 7 | 58.33% | 0 | | 0 | | 0 | |
| | 計 | 17 | 6 | 35.29% | 1 | 5.88% | 10 | 58.82% | 0 | | 0 | | 0 | |
| 40 ～ 74 | 男 | 29 | 3 | 10.34% | 1 | 3.45% | 23 | 79.31% | 0 | | 2 | 6.90% | 0 | |
| | 女 | 31 | 6 | 19.35% | 0 | | 24 | 77.42% | 0 | | 1 | 3.23% | 0 | |
| | 計 | 60 | 9 | 15.00% | 1 | 1.67% | 47 | 78.33% | 0 | | 3 | 5.00% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 77 | 15 | 19.48% | 2 | 2.60% | 57 | 74.03% | 0 | | 3 | 3.90% | 0 | |

22. 乳房マンモグラフィ検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 138 | 120 | 86.96% | 0 | | 14 | 10.14% | 0 | | 4 | 2.90% | 0 | |
| | 計 | 138 | 120 | 86.96% | 0 | | 14 | 10.14% | 0 | | 4 | 2.90% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 232 | 195 | 84.05% | 0 | | 24 | 10.34% | 0 | | 13 | 5.60% | 0 | |
| | 計 | 232 | 195 | 84.05% | 0 | | 24 | 10.34% | 0 | | 13 | 5.60% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 1 | 1 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 371 | 316 | 85.18% | 0 | | 38 | 10.24% | 0 | | 17 | 4.58% | 0 | |

23. 乳房超音波検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|--------|------|----|---------------|--------|-----|----|-------|-------|-----|----|
| | | | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 | 人数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 245 | 155 | 63.27% | 0 | | 83 | 33.88% | 0 | | 7 | 2.86% | 0 | |
| | 計 | 245 | 155 | 63.27% | 0 | | 83 | 33.88% | 0 | | 7 | 2.86% | 0 | |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 246 | 121 | 49.19% | 0 | | 116 | 47.15% | 0 | | 9 | 3.66% | 0 | |
| | 計 | 246 | 121 | 49.19% | 0 | | 116 | 47.15% | 0 | | 9 | 3.66% | 0 | |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 3 | 2 | 66.67% | 0 | | 1 | 33.33% | 0 | | 0 | | 0 | |
| | 計 | 3 | 2 | 66.67% | 0 | | 1 | 33.33% | 0 | | 0 | | 0 | |
| 合計 | | 494 | 278 | 56.28% | 0 | | 200 | 40.49% | 0 | | 16 | 3.24% | 0 | |

24. 婦人科検査

| 年代 | 性別 | 受診者数 | 正常範囲 | | 軽度異常 | | 要経過観察 要再検査 | | 要治療 | | 要精密検査 | | 治療中 | |
|---------|----|------|------|---------|------|-------|---------------|-------|-----|----|-------|-------|-----|-------|
| | | | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 | 数 | 割合 |
| ～ 39 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 291 | 282 | 96.91% | 0 | | 5 | 1.72% | 0 | | 2 | 0.69% | 2 | 0.69% |
| | 計 | 291 | 282 | 96.91% | 0 | | 5 | 1.72% | 0 | | 2 | 0.69% | 2 | 0.69% |
| 40 ～ 74 | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 715 | 690 | 96.50% | 6 | 0.84% | 13 | 1.82% | 0 | | 5 | 0.70% | 1 | 0.14% |
| | 計 | 715 | 690 | 96.50% | 6 | 0.84% | 13 | 1.82% | 0 | | 5 | 0.70% | 1 | 0.14% |
| 75 ～ | 男 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 女 | 26 | 26 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| | 計 | 26 | 26 | 100.00% | 0 | | 0 | | 0 | | 0 | | 0 | |
| 合計 | | 1032 | 998 | 96.71% | 6 | 0.58% | 18 | 1.74% | 0 | | 7 | 0.68% | 3 | 0.29% |